DOCUMENT RESUME

ED 040 546 EC 005 947

AUTHOR	Chalfant, James C.; Scheffelin, Margaret A.
TITLE	Central Processing Dysfunctions in Children: A Review of Research.
INSTITUTION	Illinois Univ., Urbana. Inst. of Research for Exceptional Children.
SPONS AGENCY	National Inst. of Neurological Diseases and Stroke (DHEW), Bethesda, Md.
REPORT NO	NINDS-Monogr-9
PUB DATE	69
NOTE	150p.
AVAILABLE FROM	Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402 ($1.25)
EDRS PRICE	EDRS Price MF-$0.75 HC Not Available from EDRS.
DESCRIPTORS	Auditory Perception, *Exceptional Child Research, Haptic Perception, Information Processing, Language Development, *Learning Disabilities, Learning Processes, *Medical Research, Memory, *Neurological Defects, Research Needs, *Research Reviews (Publications), Sensory Integration, Symbolic Learning, Visual Perception, Written Language

ABSTRACT

Research on central processing dysfunctions in children is reviewed in three major areas. The first, dysfunctions in the analysis of sensory information, includes auditory, visual, and haptic processing. The second, dysfunction in the synthesis of sensory information, covers multiple stimulus integration and short-term memory. The third area of research, dysfunctions in symbolic operations, concerns auditory language, decoding and encoding written language, and quantitative language. In addition, research needs are summarized. (JD)

PRINTED BY THE STANDARD REGISTER COMPANY, U. S. A.

FORM 0801

EDO 40546

NINDS MONOGRAPH No. 9

CENTRAL PROCESSING DYSFUNCTIONS IN CHILDREN:

A Review of Research

EC 05 487

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service National Institutes of Health

ED040546

NINDS MONOGRAPH No. 9

CENTRAL PROCESSING DYSFUNCTIONS IN CHILDREN:

A Review of Research

Phase Three of a Three-Phase Project

by

James C. Chalfant, Ed. D.
and
Margaret A. Scheffelin, Ph. D.

Institute for Research on Exceptional Children
University of Illinois
Urbana, Ill.

This report is based on research performed
under Contract No. PH–43–67–61

with the

National Institute of Neurological Diseases and Stroke
Public Health Service National Institutes of Health
U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Bethesda, Md. 20014

1969

U.S. DEPARTMENT OF HEALTH, EDUCATION & WELFARE
OFFICE OF EDUCATION

THIS DOCUMENT HAS BEEN REPRODUCED EXACTLY AS RECEIVED FROM THE PERSON OR ORGANIZATION ORIGINATING IT. POINTS OF VIEW OR OPINIONS STATED DO NOT NECESSARILY REPRESENT OFFICIAL OFFICE OF EDUCATION POSITION OR POLICY.

NINDS Monograph Series

This publication is one of a series containing neurological science contributions of NINDS staff and consultants. Included are reviews, reports, nomenclature, classification and methodology aids, and proceedings of meetings sponsored by the Institute.

Single copies of NINDS monographs are available on request.

Information Office
National Institute of Neurological
Diseases and Stroke

For sale by the Superintendent of Documents, U.S. Government Printing Office
Washington, D.C. 20402 - Price $1.25 (paper cover)

PREFACE

The increasing complexity of our modern society has created increased pressures on each individual for a greater degree of conformity to expected norms of ability and performance. These expectations have their heaviest influence during the school years, for it is at this time that certain arbitrary standards of achievement are most rigidly applied, and that the individual is subjected to the most rigid and intensive comparative evaluation. Within this system, it has long been recognized that many children do not measure up to expectations. The intelligence test was developed as a measure or predictor of academic ability, and the concept of mental retardation reflects a widespread acceptance of the fact that there are quantitative differences in the overall intellectual abilities of children.

The fact that there may be significant qualitative differences in the abilities of children has been far slower in receiving adequate recognition. The last few years, however, have seen an increasing awareness that many children, whose overall intelligence appears normal, still exhibit peculiarities or deficiencies of their mental processes which interfere with their ability to cope with certain of the standard educational requirements. With special instruction, appropriate to the specific characteristics of the individual child, many are able to surmount those difficulties and move ahead to normal or superior academic and social achievement. As the existence and the needs of this group of children have become more apparent, increasing concern has developed for the more precise definition of the problem; the delineation of the programs and services required to meet it, and the evaluation of our present state of knowledge and research.

In 1963, under the auspices of the several interested agencies of the U.S. Department of Health, Education, and Welfare, and the National Easter Seal Society for Crippled Children and Adults, a committee was assembled to consider this problem. It recommended the establishment of three task forces. The first on Definitions has completed its report.* It recommended the use of the term "minimal brain dysfunction" for a group of children of normal overall intelligence, who exhibit certain characteristics of learning or behavior attributable to a dysfunction of the nervous system. This term, which assumes all learning and behavior to be a reflection of brain function, emphasizes that at a given time it is the child rather than the environment that is different. It avoids the issue of causation, recognizing that disorder or peculiarity of brain function may stem from many causes. It avoids the more limited term "learning disability" because the disturbances of behavior, in many instances, extend further than the learning situation or the classroom.

The second task force has carried out an analysis of the medical and educational services required for children with minimal brain dysfunction. Its report is in the process of publication.

The report of Task Force III, prepared by Drs. Chalfant and Scheffelin is a review of scientific knowledge regarding the learning disabilities of these children.

---

*"Minimal Brain Dysfunction in Children," NINDB Monograph No. 3, Public Health Service Publication No. 1415. U.S. Department of Health, Education, and Welfare, 1966. (For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402. Price 20 cents.)

Its purpose is not only to summarize the current facts, but especially to point out crucial gaps in our understanding. It is a remarkable and comprehensive piece of work, highlighting above all the diversity of problems which are involved and the variety of scientific disciplines whose contributions will be required for their solution. A major problem has been the breadth of the topic and the massive literature which has been reviewed (the book includes 848 citations, but over 3,000 references are in the file). Wide gaps of knowledge exist in every area, and one is almost overwhelmed by the questions in need of elucidation.

The final summary of research needs highlights the chaotic state of our current efforts in this field. We are dealing with a poorly defined population. The methods for early recognition of the child with learning difficulties are still to be worked out and tested. There is no standard or generally accepted systematic screening program through which every child could be tested for a learning disability. The characterization of the individual deficit is on a very superficial basis, with the emphasis dependent largely upon the biases of one or another special school of thought. Remedial methods are found to rest on varied and shaky hypotheses, and have rarely been subjected to scientific evaluation even on an empirical basis.

The last few years have seen encouraging developments in these areas of research; however, one reaches the sobering conclusion that an all-out systematic research attack on the problem of the learning disabilities is long overdue.

RICHARD L. MASLAND.

TASK FORCE III—PROJECT CONSULTANTS

DOUGLAS L. CROWTHER, M.D.
Department of Neurology
University of California Medical Center
San Francisco, Calif.

RICHARD L. MASLAND, M.D.*
Chairman, Department of Neurology
College of Physicians and Surgeons, Columbia University
New York, N.Y.

JAMES W. MOSS, Ph.D.
Director, Division of Research
Bureau of Education for the Handicapped, United States Office of Education
Washington, D.C.

HELMER R. MYKLEBUST, Ed. D.
Director, Institute for Language Disorders
Northwestern University
Evanston, Ill.

RICHMOND S. PAINE, M.D.**
Children's Hospital and George Washington School of Medicine
Washington, D.C.

CHARLES R. STROTHER, Ph. D.
Director, Mental Retardation and Child Development Center
University of Washington
Seattle, Wash.

---

*Formerly Director, National Institute of Neurological Diseases and Stroke, National Institutes of Health, Bethesda, Md.

**Deceased.

CONTENTS

	Page
Preface	iii
List of Tables	ix
List of Figures	ix
1. INTRODUCTION	1
Purpose of the Report	2
Procedure for Conducting the Study	3
Limitations of This Report	3
Major Issues of Concern	4
References	5
DYSFUNCTIONS IN THE ANALYSIS OF SENSORY INFORMATION	
2. AUDITORY PROCESSING	9
The Auditory Processing Mechanism	9
Auditory Processing Tasks	11
Directions for Future Research	17
References	19
3. VISUAL PROCESSING	21
The Mechanism for Processing Visual Stimuli	21
Visual Processing Tasks	23
Directions for Future Research	35
References	35
4. HAPTIC PROCESSING	39
The Haptic Processing Mechanism	39
Kinds of Information Obtained by the Haptic System	41
Description, Assessment, and Treatment of Cutaneous Processing Dysfunctions	41
Description, Assessment, and Treatment of Kinesthetic Processing Dysfunctions	43
Directions for Future Research	47
References	47
DYSFUNCTIONS IN THE SYNTHESIS OF SENSORY INFORMATION	
5. MULTIPLE STIMULUS INTEGRATION	51
What is Stimulus Integration?	51
How Do Integrative Systems Develop?	52
What Are the Correlates to Stimulus Integration?	54
How Can Stimulus Integration Be Assessed?	56
What Factors Should Be Considered in Developing Remedial Procedures?	57
Directions for Future Research	59
References	59

Page

6. SHORT-TERM MEMORY 61
The Storage and Retrieval Process 62
Assessment 64
Disorders of Storage and Retrieval 65
Improving Memory 67
Directions for Future Research 69
References 70

DYSFUNCTIONS IN SYMBOLIC OPERATIONS

7. AUDITORY LANGUAGE 75
Acquiring the Auditory Language Code 75
Brain Mechanisms Underlying the Language Function 78
Description, Assessment, and Treatment of Auditory Language Dysfunctions 80
Directions for Future Research 87
References 88
8. DECODING WRITTEN LANGUAGE 91
The Reading Act 91
The Elements of the Graphic Code 92
Correlates to Reading Failure 94
Assessment and Treatment of Reading Failure 98
Directions for Future Research 105
References 107
9. ENCODING WRITTEN LANGUAGE 111
The Writing Act 111
Correlates to Writing Disorders 111
Disorders in Encoding Graphic Symbols 112
Assessment and Treatment of Writing Disorders 114
Directions for Future Research 117
References 117
10. QUANTITATIVE LANGUAGE 119
Correlates to Quantitative Thinking 119
Assessment 124
Training 127
Directions for Future Research 129
References 129

SUMMARY OF RESEARCH NEEDS

11. DIRECTIONS FOR FUTURE RESEARCH 135
Focus and Direction 135
Screening, Identification, and Referral 135
Assessment and Diagnosis 137
Intervention 141
Mobilizing Community Resources 144
The Need for Basic Research 146
A Final Recommendation 146
References 146
APPENDIX A—Definitions of a Learning Disability 147

LIST OF TABLES

		Page
1.	Emphases of Medicine, Psychology, and Education	2
2.	Auditory Processing Tasks	12
3.	Selection of Stimuli	14
4.	Mode of Response	14
5.	Steps in the Processing of Visual Stimuli	21
6.	Intrasensory Integrative Systems	51
7.	Intersensory Integrative Systems	52
8.	Significant Variables Which Should Be Considered Under SOR	57
9.	Acquiring Auditory Receptive Language: A Task Analysis	77
10.	Acquiring Expressive Auditory Language: A Task Analysis	77
11.	Acquiring Vocal-Motor Production: A Task Analysis	78
12.	A Whole-Word System of Reading: A Task Analysis	92
13.	A Sound-Symbol System of Reading: A Task Analysis	92
14.	Encoding Graphic-Language Symbols: A Task Analysis	111
15.	Developmental Hierarchy of Writing Tasks	112

LIST OF FIGURES

1.	Computer Model for Information Processing	3
2.	The Transduction of Auditory Stimuli	10
3.	The Transduction of Visible Stimuli	22
4.	The Haptic System	40
5.	The Psychoneurosensory Processes Involved in Facility With the Written Word	113
6.	Flow Chart for the Identification, Assessment, and Referral of Children Who Have Difficulty in School	136

To our teachers

Samuel A. Kirk
Professor Emeritus
University of Illinois

and

Severina Nelson
Professor Emeritus
University of Illinois

CHAPTER 1

INTRODUCTION

Public schools are confronted with the problem of providing educational programs for all children. In order to accomplish this task, school curricula are usually organized in a developmental sequence. Most children are introduced to a curriculum in nursery school, kindergarten, or first grade, and progress from one level to the next until their formal education is terminated.

Unfortunately, many children do not progress through the normal curricular sequences. When a child experiences difficulty in learning, the teacher, the building principal and the superintendent of schools are responsible for initiating action to: (*a*) Specify the behavioral symptoms which are related to failure in school-related tasks; (*b*) investigate possible causal factors; and (*c*) make recommendations for correcting, ameliorating, or compensating for poor performance.

In some cases, the cause of the learning problem may be obvious, as in blindness, severe mental retardation, emotional disturbance, deafness, or crippling and other health impairments. In other cases, the cause of the problem may not be so obvious. There are children with major learning problems who are not mentally retarded, deaf, blind, or emotionaly disturbed, nor do they have demonstrable brain damage. It has been hypothesized that these children have learning problems related to minimal brain dysfunction. Unfortunately, the traditional categories of special education programs do not provide services for these children. Without competent screening programs and diagnostic services, it is difficult to distinguish between children who have a minimal brain dysfunction, disorders or delayed development resulting from mental retardation, sensory deprivation, cultural deprivation, or instructional factors. Unless these differentiations are made, a child may be placed in an educational program which may not be appropriate for his educational needs.

A review of the literature reveals that a number of different terms are being used to refer to the population in question. Among these are terms such as "brain-injured," "learning disabilities," "learning disorders," "psychoneurological learning disorders," "developmental imbalances," and "minimal brain dysfunction syndrome." Definitions for these terms have been advanced by a number of individuals, professional organizations, and special committees. (See app. A.) It should be noted that these definitions focus on one or any combination of several biological and psychological events. These are outlined as follows:

1. Biological events
   a. Genetic events
   b. Neurophysiological events
   c. Structure and function of the nervous system
2. Psychological events
   a. Cognitive development
   b. Social-motivational development
   c. Present behavioral symptoms
   d. Educational consequences

A variety of descriptive characteristics may be found among these definitions. Characteristics which are often mentioned include disorders in one or more of the processes of thinking, conceptualization, learning, memory, speech, language, attention, perception, emotional behavior, neuromuscular or motor coordination, reading, writing, arithmetic, discrepancies between intellectual achievement potential and achievement level, and developmental disparity in the psychological processes related to education.

Some definitions, but not all, include references to etiological correlates such as: Injury or infection of the brain before, during, or after birth; genetic variations; biochemical irregularities; illness during the development of the central nervous system; and unknown causes. Definitions sometimes include statements about what the disorders are not. For example, children who are mentally retarded, sensorily impaired, culturally deprived, poorly instructed, or emotionally disturbed are often excluded by definition.

Table 1.—Emphases of Medicine, Psychology, and Education

Discipline	Levels of assessment	Relevant factors	Definitional emphasis
Medicine	Physiological	Etiology Prevention: Medical treatment: Changes in function and structure:	Biological events: Genetic events: Neurophysiological events: Structure and function.
Psychology	Psycho-educational correlates to learning:	Measurement Cognitive development: Remedial treatment:	Cognitive development: Pyschological events:
Education	Behavior	Prevalence Classification: Management: Behavior modification: Developmental methods: Corrective treatment:	Educational consequences: Social-emotional behavior: Motivation: Observable behavior:

Differences in terminology and definition should not be surprising since the problem area spans different disciplines including special education, psychology, speech correction, child development, neurology, and medicine. In some instances, definitions have relevance for administrative classification, while other definitions have relevance for developmental change in function and structure, adaptive behavior, diagnosis, remediation, or preventive measures. When professionals with different backgrounds, theoretical orientations, and varied professional responsibilities focus on the same problem area, differences in terminology and definition are inevitable. (See table 1.)

To clarify the problems of terminology and identification of children who have deviations in nervous system function, the National Institute of Neurological Diseases and Stroke, the National Society for Crippled Children and Adults Inc., and the Easter Seal Research Foundation provided the initiative for a special task force to study the problems of terminology and identification. The result of this task force was an extensive list of symptoms attributed to children with minimal brain dysfunction which further emphasized the behavioral diversity of the population in question (Clements, 1966). In order to provide a common term and definition which would have relevance for as many professional and lay groups as possible, the term "minimal brain dysfunction syndrome" was selected by this task force (see app. A).

The term "dysfunction" is not intended to imply etiology and was selected to avoid the implication that all individuals with this group of symptoms necessarily have demonstrable brain damage. Among the children classified as having minimal brain dysfunction are those with clear-cut organic injury of the brain. A large number of children, however, do not have a lesion in the strictest sense of the word. The learning problem may stem from a constitutional deviation in the development of the central processing system and remain undiagnosed because the neuropathological techniques for diagnosis sometimes fail to identify those deviations. Perhaps the greatest value of the term "minimal brain dysfunction" is the implication that the problem arises from within the child and not from environmental factors.

As Gallagher (1966) suggests, the criteria for selecting a definition should be one of relevance. It may be unrealistic to expect a single definition to hold relevance for many different disciplines. That which is relevant for the neurologist and pediatrician, for example, may not be relevant for the educator. It may be necessary, therefore, to formulate several kinds of definitions, each of which would have a relevance and a function for the respective user.

Since the educator must deal with behavior, it may be helpful for the educator to use terms which are descriptive of these behavioral deviations. This report focuses attention on the deviant behaviors which arise from dysfunctions of the central processing mechanisms. More specifically, this report attempts to summarize the present status of knowledge and identify future research needs with respect to the analysis, storage, synthesis, and the symbolic use of information.

Purpose of the Report

The purpose of this report is to summarize both the present status of knowledge in and the future re-

search needs for identifying, assessing and treating children with central processing disorders arising from minimal brain dysfunctions. A number of related events led to this report. First, parents and professional groups have indicated a growing concern for children with minimal brain dysfunctions. Second, this concern has not only fostered increased interdisciplinary collaboration, but has intensified conflicts between professional groups. Third, the public schools have been developing a variety of diverse educational programs for meeting the educational needs of these children. Fourth, the efforts of the public schools are frustrated by the lack of effective procedures for identification, assessment, and treatment. Fifth, institutions of higher education are beginning to mobilize their resources for training needed personnel, but there seem to be differences of opinion among school districts and State offices of education about the requirements necessary for preparing qualified teachers.

Recognizing these problems, the National Institute of Neurological Diseases and Stroke awarded a 2-year contract to the University of Illinois for the purpose of gathering information on the status of research on minimal brain dysfunction in children. The scope of work included:

(a) Exploration of the field of research pertinent to the assessment and treatment of minimal brain dysfunction in children.
(b) Delineation of the special areas which are significant for the purpose of this study.
(c) Consultation with specialists in the various fields of research pertaining to brain dysfunction.
(d) Review of the literature to identify significant studies undertaken and the researchers involved.
(e) Site visits to the various centers where pertinent research is being conducted.
(f) Compilation of information and research data and/or case histories on the status of research among the various research teams, their accomplishments and approaches taken in the assessment and treatment of minimal brain dysfunction.
(g) Preparation of narrative reports on the findings of site visits supported by bibliographic, case history, and research data materials, as applicable.

Procedure for Conducting the Study

The first 9 months of the project were spent in determining the scope of the problem, reviewing the literature and gathering pertinent information. During



Figure 1.—Computer Model for Information Processing

the data-gathering phase, a number of site visits were made to various centers to consult with specialists in the various fields of research. Journals, periodicals, and books were reviewed to identify the different problem areas which were pertinent to the purpose of the study. Research studies from education, psychology, child development, biology, biochemistry, neurology, and many other disciplines all seemed to have some relevance to the problem.

Approximately 5,000 references were identified screened, and categorized. Abstracts were made of the more significant studies. It is obvious that the large number of studies and the time limitation make it impossible to cite all the significant studies. An effort has been made, however, to include representative studies as the basis of the report.

In order to communicate with the reader it was necessary to develop a conceptual framework into which the research could be placed. The computer model for information processing was adapted for this purpose (see Fig. 1). Auditory, visual, and haptic stimuli (or sensory information) are transmitted to the central processing mechanism (brain) where they are analyzed, integrated and stored. The behavioral response of the subject serves as an additional input source (feedback) for correcting or adjusting further behavioral responses. This model was used as the framework for categorizing and relating research studies from different disciplines. The remaining 15 months were spent outlining chapters, organizing the key references, integrating information, and writing the report.

Limitations of This Report

In view of the large mass of literature which has been written about the problem area, it was necessary to

develop guidelines for delimiting the scope of the project so it could be completed during the 2-year duration of the study.

The first guideline was to focus on behavioral problems which have particular relevance for educators.

The second guideline was to report research on children, as much as possible. It should be noted, however, that in the absence of research on children, studies on adult subjects and animals have been included.

The third guideline was to report research and generally exclude literature which reflected opinion. In the event that research data were not available, unsupported theoretical hypotheses occasionally have been included.

The fourth guideline was to exclude the large mass of research with respect to emotional disturbance and social maladjustment. While these behavioral deviations are most certainly relevant to education, they represent a large and distinct problem area which merits special study in its own right. Some mention of these problems is included in this report, however, to indicate how they relate to the particular content area under discussion.

Major Issues of Concern

In reviewing the research, three major bodies of literature were identified. First, research studies from experimental child psychology typically report findings on the responses of normal children to discrete auditory, visual, or haptic qualities of stimuli. Second, research studies from clinical psychology and clinical education report findings on the responses of atypical children with perceptual disorders to the objects themselves. Third, research studies from the medical profession report findings on neurophysiological correlates. The experimental child psychologist has focused his attention on children's responses to discrete stimulus qualities such as shape, color, form, size, area, etc., while the clinical educator and the clinical psychologist have been attempting to teach children to respond to and manipulate objects which represent multiple stimulus qualities. Unfortunately, the findings from experimental research have not been translated into useful procedures for the clinical assessment and training of children, nor have the clinical procedures been described in the specific details necessary for replication and experimentation. Most of the research by the medical profession on organic brain damage has been conducted with adult subjects. Comparatively little information is available about children who have suffered known brain damage.

As a result of these different approaches and independent research efforts, there are gaps in the present status of knowledge for those who attempt to relate the research findings of experimental child psychology, clinical education, and clinical psychology, and the medical profession. One of the major objectives of this report is to attempt to relate the research findings from these three disciplines.

A number of major issues with respect to the identification, assessment, and treatment of central processing dysfunctions were highlighted by the review of the literature. Some of the more important issues are presented here in question form and are discussed throughout the report.

1. What are central processing tasks?
2. What is the significance of processing tasks for learning?
3. What are the anatomical, neurological, and physiological components which constitute the central processing mechanisms?
4. What observable behaviors are symptomatic of central processing dysfunctions?
5. How are these behavioral symptoms linked with organic brain dysfunction?
6. How effective are the procedures for identifying and assessing the severity and extent of central processing dysfunctions?
7. What aspects of diagnosis are most relevant for educational intervention?
8. How effective are the procedures for preventing, ameliorating, or compensating for specific central processing dysfunctions?
9. How can research design be sharpened to determine the effectiveness of specific treatment procedures for specific kinds of dysfunctions?
10. What future research is needed to resolve these issues?
11. Where does the responsibility lie for conducting needed research?
12. How can research findings be disseminated to the practitioner in the field?

To make an authoritative analysis of this problem, the authors should be highly specialized in anatomy, neurology, biochemistry, physiology, audiology, vision, experimental child psychology, educational psychology, learning theory, psycholinguistics, and behavior modification, all of which are related to central processing dysfunctions in children. It is obvious that no author is a specialist in all these areas. In our particular case we were fortunate enough to have consultants from

these areas to assist us and to criticize the manuscript. The authors may be more accurately described as consumers of research. The difficulty we encountered in attempting to pull together and integrate the mass of conflicting and sometimes confusing information may be typical of the difficulty experienced by other professionals. At best, this report represents an ordinary effort. A beginning. A search for the right questions.

REFERENCES

CLEMENTS, S. D. *Minimal brain dysfunction in children.* Public Health Service Publication No. 1415, January 1966, Washington, D.C., U.S. Department of Health, Education, and Welfare.

GALLAGHER, J. J. Children with developmental imbalances: A psychoeducational definition. In W. M. Cruickshank (ed.). *The teacher of brain-injured children.* Syracuse, N.Y.: Syracuse University Press, 1966. pp. 23–43.

DYSFUNCTIONS IN THE ANALYSIS OF SENSORY INFORMATION

361-752 O—70——2

6/7

CHAPTER 2

AUDITORY PROCESSING

The auditory channel is one of the most important avenues through which children and adults receive information about their environment. The importance of hearing acuity for obtaining such information has been established, but there is little known about the central processing of auditory stimuli. There are children, for example, whose hearing acuity is within the normal range of hearing, but who have difficulty processing and obtaining meaning from auditory stimuli. A child who has difficulty processing auditory stimuli may be observed to perform poorly in some of the following tasks: (*a*) Identify the source of sounds; (*b*) discriminate among sounds or words; (*c*) reproduce pitch, rhythm, and melody; (*d*) select significant from insignificant stimuli; (*e*) combine speech sounds into words; or (*f*) understand the meaning of environmental sounds in general.

Clinical case studies typically have reported such observable behaviors as being characteristic of children who have difficulty in processing and responding to auditory stimuli. In many instances, the clinical report simply mentions the existence of these behaviors and does not provide a comprehensive description of the conditions under which the subject was unable to process and respond to the auditory stimuli in an appropriate manner. There is need to describe disorders in processing and utilizing auditory stimuli in more detail. The use of more precise terms will help facilitate communication between different disciplines.

Many of the observable behaviors which seem to be characterized by difficulty in perceiving auditory stimuli have been referred to as "auditory perceptual disorders." Myklebust (1954) defines auditory perception as the ability to "* * * structure the auditory world and select those sounds which are immediately pertinent to adjustment (p. 158)." According to Berry and Eisenson (1956), children with auditory perceptual disorders can hear sounds, but are unable to recognize the sounds that they hear. The term "auditory perception," as it is used here, refers to the central processing of auditory stimuli.

In 1954, Myklebust made a number of important distinctions concerning auditory disorders in children. Unfortunately, the present status of knowledge has not advanced very far beyond the early contributions of Myklebust (1954), Goldstein (1948), Nielsen (1948), and others.

Several factors may have contributed to the lack of empirical data on auditory stimulus processing, especially in children. First, is the lack of data on the nature of auditory stimuli, especially speech sounds. Second, it is difficult to measure and study responses to auditory stimuli. Third, the organization, structure and use of sound in the environment is achieved at different ages by different individuals. Fourth, confusion is generated by overlapping terms. Some terms such as "psychic deafness" refer to diagnosis. Terms such as "sound localization" refer to a specific task, while terms such as "hi-frequency loss" refer to aspects of the stimuli.

The purpose of this chapter is to provide an overview of the disorders which can occur in processing auditory stimuli. More specifically this chapter will discuss: (*a*) The processing mechanism; (*b*) different kinds of auditory processing tasks; and (*c*) the needs for future research in assessment and training.

The Auditory Processing Mechanism

A deficit in hearing acuity which blocks incoming acoustic stimuli will interfere with the perceptual process (Myklebust, 1957). For this reason, any discussion of the mechanism for processing auditory stimuli should include a description of the human ear as a transducer of acoustic stimuli, as well as a brief discussion of the cortex as an auditory processor.

The Ear as a Transducer

O'Neill (1964) has described the sequence of physical pressure changes by which acoustic stimuli are transmitted to the cerebral cortex via acoustic, mechanical hydraulic, and electrical signals. O'Neill's description of what happens to the pressure changes



Figure 2.—The Transduction of Auditory Stimuli

created by the original sound source has been adapted and expanded in figure 2. A vibrator (larynx, bell, reed) produces an initiating stimulus. The sound produced in the medium outside the ear (usually air) is transduced through the inner ear fluid, to the cochlear nerve, and then to the auditory cortex of the brain. The amount of energy that appears to be necessary for detection of a sound stimulus is extremely minute. O'Neill says a sound pressure of 0.0002 dynes per centimeter "* * * is considered the lowest intensity of sound that the human ear can detect" (p. 10). This amount is one ten-trillionth times as weak as the most intense sound that can be tolerated by the ear. "The human ear is able to detect some 340,000 * * * differences in frequency and intensity (O'Neill, 1964, p. 20)."

The Cortex as an Auditory Processor

The auditory cortex has been viewed as a passive receptor of auditory stimuli which are transmitted along the auditory nerve. More recently, it has been conceptualized as a selective analyzer. According to the theory of selective-analysis, "* * * sensation incorporates the process of analysis and synthesis of signals while they are still in the first stages of arrival (Luria, 1966, p. 97)." This may result in increased excitability and sensitivity with respect to some components of the stimulus such as color or in decreased excitability with respect to others such as form (Granit, 1955; Sokolov, 1958).

According to this theory, the sensory divisions of the cortex seem to be responsible for the analysis and integration of complex signals and are not responsible for the first stage, reception of sensation. Therefore, lesions in this area will not necessarily alter the threshold of sensation, but will instead result in a disturbance of the higher, analytic-synthetic function. Research workers in both the Soviet Union and in the United States have found that extirpation of the sensory cortex in animals disturbs the ability to differentiate between pairs of both simple and complex signals, and results in an incapacity to form new conditioned reflexes in response to complex sound stimuli (Kryzhanovskii, 1909; Babkin, 1910; Butler, Diamond and Neff, 1957; Goldberg, Diamond, and Neff, 1957). Findings such as these suggest that pathology of the auditory divisions of the cortex is reflected in the more complex forms of differential auditory analysis and not in hearing acuity (Luria, 1966).

The reflex theory of higher mental functions including auditory analysis rejects the localization of sensory perception in an isolated area of the brain. The reflex theory, which views mental functions as complex reflex processes having their origin in the child's response to adult verbal commands, regards the brain as a system of acquired complex intercentral connections (Luria, 1966). While this theory of auditory analysis may give us a more substantial basis for understanding the behavioral symptoms which arise from lesions in the auditory divisions of the cortex, it is very difficult to

study, in detail, the role played by the cerebral cortex in auditory functions.

There is some evidence which suggests: (*a*) That the primary receiving strip for auditory stimuli is located in the Sylvian margin of the temporal lobe, area 41 (Penfield and Rasmussen, 1950); and (*b*) that the secondary division of the first temporal convolution is responsible for the analysis and integration of sound signals (Luria, 1966). It should be noted that most of the research conducted with human subjects has been with adults who have suffered tumors, abscesses, or traumas to the temporal lobe.

As early as 1873, Ferrier (1876) demonstrated in animals the localization of the auditory center in the temporal lobe. Electrical stimulation of the temporal lobe of either side close to the fissure of Sylvius will produce auditory sensations. Campbell (1905) recorded the location of points which, when stimulated, appeared to produce auditory sensations in humans. A variety of auditory sensations in the contralateral ear or in both ears were recorded. When the cortex was stimulated in the same place in different patients, different sensations were reported. Thus, the lack of interpatient reliability makes it difficult to analyze in detail the precise function of the cortex in auditory functioning.

The research concerning the function of the secondary divisions of the auditory cortex is less clear. It may be here that the complex forms of auditory analysis and integration occur (Penfield and Rasmussen, 1950; Luria, 1966). This region may be largely responsible for the systematic deciphering of sound signals necessary for the perception of speech. According to Luria (1966):

> * * * the disturbance of discriminative hearing, which can now be interpreted as a disturbance in the analytic-synthetic activity of the auditory cortex * * * may be regarded as the fundamental symptom of a lesion of the superior temporal region of the left hemisphere, and the resulting acoustic agnosia may be regarded as the fundamental source of speech disturbance (pp. 106–107).

It is suggested that future research should attempt to identify the fundamental defects resulting from lesions of the secondary divisions of the auditory cortex. There is need, also, to conduct research on the disturbances which arise from lesions of the middle segments of the convex portion of the left temporal region. This area has a high incidence of lesions (tumors and abscesses) and the disturbances associated with these lesions have not been clearly defined. The clinical behaviors are varied and complex, and include auditory and visual hallucinations, dreamy states, emotional changes, and changes in the state of consciousness (Penfield and Roberts, 1959).

Auditory Processing Tasks

The term "auditory perception" has been used to describe many behavioral responses to auditory stimuli. In order to present an organized view of auditory perception or auditory stimulus processing, seven different auditory tasks have been identified and are presented in table 2. It will be noted that the seven tasks are described according to the stimuli presented, the response required, and terms commonly applied either to the task or to failure in performing the task. Each of the seven tasks will be described in separate sections, which will include whenever possible a discussion of the: (*a*) Nature of the task; (*b*) neurophysiological correlates; (*c*) consequences of failure in performing the task; (*d*) remedial procedures; and (*e*) future research needs.

1. *Attention to Auditory Stimuli.* The first concern in testing or teaching responses to auditory stimuli is to determine whether or not the child is attending to the stimuli which are being presented. Inattentiveness to auditory stimuli might be related to: (*a*) Low level or absence of hearing acuity; (*b*) distractibility involving competitive visual or auditory stimuli; (*c*) hyperactive behavior; (*d*) severe emotional disturbance; (*e*) severe mental retardation; or (*f*) inability to obtain meaning from auditory stimuli.

A thorough differential assessment of children who seem to have difficulty processing auditory stimuli should include an examination of all the correlates to the attentional factor. It is important to remember that attention to auditory stimuli is inferred from the subject's responses such as inclining one's head toward the source of sound, facial expressions, or verbal or motor responses. There is need to develop systematic procedures for assessing the reasons for what appears to be inattentiveness to auditory stimuli.

There is little research available about the most efficient ways to teach the child to attend to auditory stimuli. The literature typically describes clinical suggestions or approaches to this problem. For example, the use of amplification can help intensify auditory stimuli and create an awareness of the differences between sound and no sound. Toys, musical instruments, and household appliances which can be manipulated by the child can be used for training purposes. Suggestions include repeatedly turning the child's head toward the source of sound, or making the sound source visible to the child when he turns his head.

Table 2.—Auditory Processing Tasks

Stimulus presented	Response required	Common terminology
1: Auditory stimulus	Indicate awareness through verbal or motor response:	Attentional problem, distractible, hyperactive:
2: Sound versus no sound	Yes/no	Acuity, detection:
3: Sound from several different origins	Indicate direction from which originated	Sound localization:
4: Sounds varying on one acoustic dimension:	Same/different	Discrimination of pitch, loudness, speech sounds, noises:
5: Sequences and patterns of speech or nonspeech sounds varying on more than one acoustic dimension:	Reproduce sequence: (a) Imitation, e.g:, tapping; (b) speaking; (c) singing:	Pitch, rhythm, melody, "arhythmical, can't carry a tune, tone-deaf, poor auditory memory:"
6: Sound preselected as "figure" versus sound preselected as "ground."	Select "figure" sound	Differentiate, discriminate.
7: Sounds from one or more sources	Identify by: (*a*) Pointing to a visual representation of the sound source; or (*b*) naming the sound source:	Associating sounds with their actual sources:

Selective reinforcement, contingent upon response, and other principles of behavior modification can be used to help increase attentiveness to auditory stimuli (Sloane and MacAulay, 1968). There is need to conduct research on the ways in which attentiveness to auditory stimuli can be increased in those individuals who tend to ignore the significance or meaning of sound.

2. *Sound Versus No Sound.* The first level of assessment of sound discrimination is to indicate whether or not the sound has been heard. A basic procedure in assessing sensitivity to sound, for example, is to present pure tone or warbled tone stimuli to determine if the child can differentiate sound from no sound. Lowered acuity in hearing alone does not represent an auditory perceptual disorder. For auditory processing to occur, however, the hearing mechanism must be capable of transmitting the mechanical, hydraulic, and electrical signals to the auditory cortex. A hearing deficit will reduce the accuracy of discrimination.

3. *Sound Localization.* Some children have difficulty in localizing or indicating the source or the direction of a sound. When the source of sound is equidistant from both ears, the sound is said to be difficult to locate even by practiced adults. If the sound source is moved, either to the right or to the left of the midline of the body, it is closer to one ear than the other, and the closer ear is stimulated somewhat earlier. Since the acoustical complexity of a sound is partially a function of distance from the ear, the ear closest to the sound will receive a stimulus of greater complexity than the ear farther away. The sound should then appear different to the two ears. The time difference between the detection of the signal presented to each ear should also be a clue which can be used to detect the source of sound. Children who have difficulty in identifying the source of sound may not learn that different people have different voices or that the sound made by one person is specific to that person and not produced by a random source.

In reviewing the literature, no data were found on the assessment or training of sound localization. There is need to develop activities which will improve the ability to localize sounds in terms of distance as well as direction. Being able to localize sounds will help the person visually link sounds with their sources and help establish associations between sounds and objects or events.

4. *Discriminating Sounds Varying on One Acoustic Dimension.* For our purposes, auditory discrimination is defined as indicating whether two acoustic stimuli are the same or different. Given a pair of auditory stimuli, the subject must indicate whether the two members of the pair are alike or different. The response required varies from vocal ("same-different," "same-not the same," "now I hear it") to various forms of nonverbal communication (turning toward the source of sound, performing an agreed upon action representing "same" or "different," pointing to a pair of similar objects, rather than to a pair of unlike objects).

Auditory stimuli may vary along several acoustic dimensions and several presentational dimensions. Individual acoustic stimuli can be measured on several physical scales such as frequency and intensity. These physical scales are related to auditory-psychological

dimensions such as pitch and loudness. Presentational dimensions which include number, rate, and duration of stimuli have auditory-psychological counterparts in rhythm and melody. The kind of sound (speech versus nonspeech sounds) and the location of the sound are also presentational dimensions. Auditory stimuli may be presented either simultaneously or successively to one ear or both ears. The relationship of the hearer to the acoustic stimulus will affect his judgment of the auditory stimulus. For example, the farther away a person is from the source of the acoustic stimulus, the lower the pitch and the softer the intensity will appear to be.

The assessment of differential responses to pairs of auditory stimuli involves discriminations along one or more acoustic or presentational dimensions. There are two major classes of sound sources. There are human sources (vocal sounds and words, etc.) and nonhuman sources (bells, watches, machines, etc.). Most standardized discrimination tasks include only a few different sound sources and may include stimuli from either one or both classes of sound sources.

In view of the fact that a large proportion of human communication consists of speech, a series of vocal acts, it is not surprising that several tests of speech-sound discrimination have been developed. Among the most widely used tests are the Wepman Auditory Discrimination Test (Wepman, 1958), the PERC test (Drake, 1965), and the Boston University Speech Sound Discrimination Picture Test (1955). Norms are available for the Wepman and are being secured for the PERC. The chief difference between the two tests is that the child is required to say "same" or "different" to each pair of words on the Wepman, whereas he is required to say "same" or "not the same" on the PERC. Extensive descriptions of several widely used speech sound discrimination tests, and lists of the words or sounds presented to the subject, are supplied by Berry and Eisenson (1956, pp. 449–501). Caution should be exercised in interpreting scores from these tests, because of the problems in presenting vocally produced stimuli in a consistent manner. It is possible that the repeated readings of the list of words do not necessarily give equivalent forms of a test because of day-to-day differences in the speaker's presentation.

Disorders in sound discrimination have been found to be caused by lesions in the temporal lobe. Some individuals experience difficulty in identifying similar sounds, which they perceive as being different (Korst and Fantalova, 1959). Difficulty in discrimination also occurs when sound complexes are presented which have the same components but are presented in a different order (Traugott, 1947; Babenkova, 1954). A disturbance of discriminative hearing may result from a lesion of the superior temporal region of the left hemisphere. Luria (1966) views a disturbance in the analytic-synthetic activity of the auditory cortex, and the resulting acoustic agnosia, as the fundamental source of speech disturbance.

There are individuals who have adequate hearing acuity, but who do not discriminate one sound from another. Failure to discriminate between auditory stimuli has a number of possible consequences. If children are unable to hear the differences or similarities in initial or final sounds of words, consonant blends, or vowels, they will have difficulty in acquiring, understanding, and using spoken language. Some individuals have difficulty in distinguishing between single speech sounds. It is more difficult to discriminate between similar sounds (/d/, /t/, /p/) than if the sounds are widely different (/h/, /k/, /s/). Some individuals are aware that a difference exists between two sounds, but may not be able to specify the nature of the difference.

The analysis and synthesis of series of speech sounds is basic to learning the phonemic structure of language. A child who is unable to analyze series of sounds into their separate parts or to synthesize separate sounds into wholes, may suffer impaired auditory memory, speech, and reading ability. Monroe (1932), for example, found that among first grade children, a group of nonreaders made significantly more errors on auditory discrimination tasks than did a group of adequate readers. Yedinack (1949) and Monroe (1932) found a high proportion of poor readers had speech defects. Monroe concluded that either the speech defect was a factor in reading disability or both the reading and speech problems were the result of a common cause, probably the inability to discriminate successfully the sounds of words. Chronological age of the child is an important consideration, because many children continue to improve in discriminating sounds as late as the eighth year (Wepman, 1960).

If we are to understand the processing of auditory stimuli, there is need to study the ways in which an individual processes auditory stimuli varying along different acoustic and presentational dimensions. In reviewing measurement techniques, Reichstein and Rosenstein (1964) recommended that four important variables should be studied. These include the selection of the stimuli, the mode of input, the method of response, and motivational factors. They have summarized both the various modes of presenting auditory stimuli and the kinds of responses that are required. The pure tone audiometer is used to present auditory

stimuli. Similar to the pure tone is the warbled pure tone which differs only in that the tone is not steady. When the auditory stimulus is speech, word lists or word games are often used. Complex nonspeech stimuli such as music, noise makers, and animal sounds are frequently employed as another mode of presenting auditory stimuli.

The child may be required to respond in a variety of ways. His perception of the auditory stimulus may be measured by a reflex response which the stimulus evokes or by an indication of locating the stimulus. The child may be required to voluntarily indicate his perception by saying something or raising his hand. In the conditioned response method the child is conditioned to reach for or do something pleasant each time he perceives an auditory stimulus. In the simple "play" conditioned response, the child performs a simple motor act when he hears the auditory stimulus. In the complex "play" conditioned response upon hearing tone the child is to respond by performing some action with a complex toy.

Lists of representative studies related to the selection of auditory stimuli and mode of response are presented in tables 3 and 4. These lists, while not exhaustive of the kinds of variables that should be accounted for, demonstrate the wide variation in the procedures employed. In addition, pretest training to teach children response alternatives is an aspect of testing that has not received sufficient attention.

There is need to develop and program sequential training activities which will help individuals discriminate the differences between sounds. The concept of auditory discrimination is rather broad. Research efforts in this area, therefore, probably should be directed toward the discrimination of different kinds of auditory stimuli. Variables such as duration, pitch, frequency, volume, symbolic or nonsymbolic stimuli, speech sounds, and noises should be considered.

Training should follow the principle of beginning with maximum contrast between the members of the pair to be discriminated. Discrimination should begin with gross sounds commonly heard as those found on training phonograph records (Utley, 1950) and proceed to finer sound differences. Mecham (1966) suggests extensive training in recognizing and discriminating the sound or sounds which are later to be reproduced. According to Berry and Eisenson (1956), some children require more time to absorb material and will respond more readily if the stimuli are intensified and repeated. Miller (1951) suggests that children can be trained in the automatic repetition of sequences such as the number series 1–10. After the child has mastered the auditory-vocal sequence, he can be trained to pick out the visual representation of the auditory number name. After the initial gross discrimination task has been performed successfully, the task can be made progressively more difficult by decreasing the range of differences.

Implications for training methods may be discovered in Mecham, Berko, Berko, and Palmer's (1966) ra-

Table 3.—Selection of Stimuli

Type of stimulus	Reference
1: Pure tone	Myklebust, 1954:
2: Warbled pure tone	Douglass, Fowler, and Ryan, 1961.
3: Speech	Keaster, 1947: Siegenthaler, Pearson, Lezak, 1954: Dale, 1962: Solomon, 1962: Wolfe and MacPherson, 1959: Sortini, 1960:
4: Music, noisemakers, animal sounds:	O'Neill, Oyer, Hillis, 1961: Streng, Fitch, Hedgecock, Phillips, and Carrell, 1958: Solomon, 1962: Whitehurst, 1961:

Table adapted from Reichstein and Rosenstein (1964):

Table 4.—Mode of Response

Type of response	Reference
1: Reflex-localization	DiCarlo and Bradley, 1961: Ewing and Ewing, 1958: Downs, 1960:
2: Direct-voluntary indication	Myklebust, 1954:
3: Conditioned response methods	Thorndike, Hilgard, and Skinner (in Hilgard, 1956): Miller and Dollard, 1941:
4: Complex-play conditioned response	Dix and Hallpike, 1947: Guilford and Haug, 1952: Wolfe and MacPherson, 1959: Sills, 1962: Meyerson and Michael, 1960:
5: Simple-play conditioned response	Ewing and Ewing, 1944: Lowell, Rushford, Hoversten and Stoner, 1956: Meyerson and Michael, 1960: Costa, Mandel, and Rapin, 1962:

Table adapted from Reichstein and Rosenstein (1964):

tionale for dividing learning behavior into three aspects: (1) Conditioning discriminative stimuli; (2) differential reinforcement of successive approximations of the desired behavior; and (3) chaining or sequencing behavioral patterns.

5. *Discriminating Sound Sequences Varying on Several Acoustic Dimensions.* A child may be able to discriminate one sound from another, yet experience great difficulty in discriminating or reproducing groups or patterns of auditory stimuli. Rhythm is the sequential pattern of several auditory stimuli in time. Processing auditory stimuli varying on several acoustic dimensions is an important factor in the acquisition of spoken language. At present, there is little information about the wider implications of disorders of rhythm, pitch, and their combination, melody. The purpose of this section is to explore some of the implications which have been mentioned in the literature.

Luria (1966) cites case studies of individuals who have suffered brain damage to the fronto-temporal region having a history of musical disorders in which the rhythm and melody functions are impaired but the pitch relationships remain unimpaired. The involvement of the short-term memory function in fronto-temporal disorders may help account for the difficulty in reproducing rhythm patterns, while pitch relationships remain unaffected. Pitch discrimination activities do not usually make severe demands upon the memory function, because one pitch is usually presented immediately after another. The investigation of the perception and reproduction of pitch relationships in musical melodies is an important aspect in studying pathology of the temporal and premotor divisions. Simple tests consist of comparing whether one pitch is higher or lower than or equal to a second pitch.

Semeritskaya (1945) found that patients with lesions in the temporal region can reproduce a rhythmic pattern if it is presented slowly and can be counted. These same patients, however, have great difficulty in reproducing rhythmic patterns which are presented rapidly. When the auditory analysis and integration functions are disturbed, patients can reproduce single rhythmic groups, but are frequently unable to repeat the rhythmic pattern over and over as in a series. These symptoms have been related to lesions of the superior left and right temporal regions. Auditory disturbances may not be so conspicuous in patients with lesions of the inferior division of the left-temporal region as are auditory disturbances resulting from lesions of the superior divisions.

There is at present little information as to how the perception of rhythm is disturbed by lesions in the motor system. There is some evidence that the evaluative role played by the motor analysis of rhythms suggests that lesions in the anterior divisions of the brain may interfere with the proper evaluation of rhythmic structures. With frontal lobe lesions there is difficulty in analyzing rhythmic structure. The tendency is to overestimate rhythmic patterns and to successively change rhythmic structures from groups of two to groups of three.

One of the important aspects in examining children is to determine the conditions under which they find it difficult to perform tests based on rhythm. Is it actually a problem in analyzing acoustical images? Is it a defect in the regulatory role of verbal instructions? Is it defective motor functioning? Is it the shift from reproducing one rhythmic structure and then another which is an indication of the mobility of nervous processes?

When the reproduction of rhythms is disturbed by lesions of the premotor region, the disturbance of the system of higher automatisms makes smooth reproduction of melodies impossible. In these cases each tap forms part of a rhythmic pattern and requires an isolated impulse. The child does not appear to develop an automatic presentation of the rhythmic pattern. He forms each tap individually. The difficulty is increased with an attempt to increase the tempo, and inhibition of the tapping becomes more difficult and superfluous taps appear. Although the child is aware that these taps are out of place, he is not always able to suppress them. Damage to the motor analyzer leads to difficulty in reproducing the accentuated rhythms. It is difficult for the subject to change from tapping one rhythmic group to tapping another. He is more apt to tap out the same sequence rather than the new ryhthm.

There is usually no difference between the reproduction of rhythms in the imitation of acoustic images and the production of rhythms from verbal instruction in patients with lesions of the premotor region. There is, however, a tendency toward preservation on the initial stimulus item under both imitation and verbal instruction conditions. Inability to correct mistakes may often persist when rhythms are reproduced from an acoustic image. This repeated failure suggests that the fundamental effect of the rhythms reproduced by patients with a frontal syndrome is associated with the disturbance of the selective aspects of the motor act, and of the regulatory influence in originating the motor act.

Generally, patients with lesions of the temporal divisions of the brain often have difficulty reproducing rhythms from an acoustic pattern, particularly when

the rhythms are presented quickly and the patient is unable to count them. The counting response is an element which can be controlled by the experimenter. If the child has the ability to count, patterns can be presented in such a way that counting is ruled out as a mediator. On the other hand, if the child is having extreme difficulty, perhaps counting could be introduced as a mediator to improve his ability in dealing with rhythm.

Quantitative aspects of the stimuli, such as the number of units in groups of acoustic signals, is an important consideration. The task can be made more complicated by manipulating the number of taps, complexity of pattern, and number of groups. The rapidity with which groups are presented in succession is an important temporal variable. Auditory tap patterns require the acoustic analysis and reproduction of the rhythmic structure. Apparently, the perceived acoustic structure must be coded as sounds corresponding to a series of consecutive motor movements, before they can be reproduced. This is a difficult task for some persons to perform.

On occasions when a rhythmic structure is presented at a slightly faster rate, or rhythmic counting aloud is forbidden, some patients are unable to evaluate the pattern. The complaint is that the taps are presented too quickly, the patient cannot keep up, and there may be a tendency to overestimate the number of taps in such series. For example, when slow taps are followed by two fast taps, the second group is often judged to contain three taps. Some patients do not benefit when a particular rhythmic group is repeated many times in succession. This kind of presentation normally enables the subject to conduct a longer comparison of one group with another. But other patients often believe that there are too many taps when they cannot identify the rhythm in which they are presented.

6. *Auditory Figure-Ground Selection.* Some children experience difficulty in selecting the relevant from the irrelevant auditory stimuli in their environment. Because most verbal communication takes place by auditory speech signals, a child who is unable to attend to speech sounds or to differentiate speech sounds from the remainder of the auditory stimuli in the environment will probably experience difficulty in learning to comprehend and in acquiring language as a communication system. Other behaviors sometimes include distractibility, short attention span, and ignoring some auditory stimuli in listening activites.

Assessment of the interference effects created by competing auditory stimuli may be accomplished by presenting a significant stimulus, and at the same time, presenting stimuli which have been declared by the examiner to be insignificant. The response required of the subject is to indicate either that he has heard the significant stimulus, or that he can reproduce the significant stimulus. Little is known about children's performances on a task of this type. There are many clinical reports of children who respond to insignificant auditory stimuli during testing or teaching situations. The so-called distractible child is an example of a person who appears to attend to the irrelevant auditory stimuli of a situation. In a classic study, Miller, Heise, and Lichten (1951) presented a series of words, together with random noise, to adult subjects. The subject's task was to repeat each word. The results showed that incorrect responses were influenced by the number of words in a set, the number of previous presentations, familiarity, and the acoustic power of the irrelevant auditory stimuli.

There is little information concerning the interference effects of presenting two or more auditory stimuli at the same time or at different points in time. The sequencing and timing of auditory stimuli may be an important variable. Interference effects may occur, when two auditory stimuli are presented at the same time, or one auditory stimulus is presented either before or after another. Delayed auditory feedback has been used to investigate interference effects. The air-conducted sound of a subject's voice is delayed electrically and presented at a specified delay interval to the subject by means of earphones. When two or more stimuli are presented at the same time, "masking" is said to be taking place. When two or more stimuli are presented at different points in time, "fatigue" is said to be taking place (Hirsh, 1952).

Little research has been done on methods or materials for teaching a child to select significant from insignificant auditory stimuli. There is need to conduct a systematic program of research in which various aspects of attention, organic conditions in the brain, effect of drugs, and nature and presentation of auditory stimuli are thoroughly explored.

7. *Associating Sounds With Sound Sources.* Some individuals seem to have difficulty establishing a correspondence between sounds and their producers. The inability to link sounds with their sources may involve correlates such as intelligence, auditory memory, the ability to localize sounds, and acoustic discrimination. The end result is difficulty in obtaining meaning from sound and acquiring the use of language as a means of receptive and expressive communication.

The term auditory agnosia refers to an impairment in an individual's recognition of sounds or combina-

tions of sounds and his attachment of meaning to those sounds. The problem is not one of acuity. The person is aware of sounds and hears sounds but does not relate these sounds to other experiences.

When the auditory agnosia is confined to speech sounds, it is described as word deafness or auditory aphasia. In some cases auditory agnosia is confined to nonspeech sounds such as mechanical or animal noises. Also, an agnosia for musical sounds, melody patterns, and rhythm has been reported. Myklebust (1957) distinguishes between auditory agnosia and aphasia as follows:

> * * * the aphasic finds all sounds in his environment useful and meaningful with the exception of the spoken word. In contrast, the auditory agnosic not only cannot use these spoken sounds in this environment but he cannot attribute meaning to any sounds in his auditory world (p. 511).

Myklebust points out:

> Severe auditory perceptual disturbances and auditory agnosia are highly similar in symptomatology but comparatively the condition of agnosia seems to be considerably more severe. It seems that an auditory agnosia does not occur unless an aphasia also is present (p. 511).

The person with an agnosia for acoustic stimuli is unable to recognize sound patterns and requires training to establish associations between sounds and situations, sounds and their sources, and sounds and actions.

In 1869 Bastian (1898) described "word deafness" in which the adult subject was able to hear, but was unable to recognize words. Wernicke (1874) hypothesized the location of the general auditory speech area of the brain in the first temporal convolution. A loss in this area was found to produce a loss in understanding speech. Nielsen (1948) reported cases in adults which show that word deafness is due to a lesion in the middle third of the first temporal convolution on the major side. Damage to this area seemed to result in failure to recognize spoken words. The adult subject does not understand anything that is said to him, nor can he repeat words, nor write from dictation, but spontaneous speech may not be affected.

In acoustic agnosia the subject is unable to recognize the sounds he hears. Auditory verbal agnosia results from a lesion of Wernicke's area and occasionally from lesions in its vicinity within the temporal lobe.

Directions for Future Research

In contrast to the body of knowledge which has been gathered on hearing acuity, comparatively little research has been done on the processing of auditory stimuli. There is need to more clearly identify auditory processing tasks, and describe and categorize the observable behaviors which are associated with these tasks. There is a need to describe the behavioral symptoms which characterize efficient auditory processing, as well as dysfunctions in auditory processing.

Research is needed to explore the behaviors of auditory processing disorders related to: (*a*) attention to auditory stimuli; (*b*) differentiating sound from no sound; (*c*) sound localization; (*d*) discriminating sounds varying on one acoustic dimension; (*e*) discriminating sound sequences varying on several dimensions; (*f*) auditory figure-ground selection; and (*g*) associating sounds with sound sources. The use of precise terms will help facilitate communication about these behaviors.

Basic research is needed in neurology and biochemistry if we are to link specific behavioral symptoms or clusters of behaviors to neurological and physiological correlates. The present status of technology and the relative inaccessibility to the human brain makes it difficult to study the role of the cortex in auditory functions. While some research has been done, there is need to improve our technology and attempt to increase our knowledge of neurophysiological behavior. Unless this happens, assessment and classification will continue to be substituted for differential diagnosis.

The lack of reliable and valid diagnostic procedures and the lack of standardized terminology make fine diagnostic differentiation a difficult task. DiCarlo (1960), for example, reevaluated 67 children who had been diagnosed as aphasic by other diagnosticians. He found that 28 children were mentally retarded, 15 peripherally deafened, and 20 were emotionally disturbed. He found only four to be aphasic. It is difficult to identify the causes of auditory processing disorders, because different etiological factors are often characterized by many of the same behavioral symptoms. Failure to respond to auditory stimuli may be attributed to peripheral deafness, central deafness, mental retardation, severe emotional disturbance, aphasia, or to auditory imperception (Myklebust, 1954). One of the most basic research steps which should be taken is to attempt to provide more detailed and comprehensive descriptions of the behavioral responses to auditory stimuli which differentiate these conditions.

There is some evidence, for example, that auditory behavior tends to be consistent with mental age and that the mentally retarded seem to respond better to meaningful test stimuli than to the more abstract pure tone test (Myklebust, 1954; Wolfe and MacPherson, 1959). There is need, however, to develop diagnostic procedures to further differentiate mental retardation

from other etiological factors. Also, children with peripheral hearing losses have been found to use their residual hearing in a consistent and meaningful manner (Myklebust, 1954; Whitehurst, 1961).

Children who give consistent, although frequently incorrect, responses to an auditory stimulus are typically classified as "peripherally deaf." On the other hand, children who respond inconsistently are typically classified as "centrally deaf," "auditory agnostic," or "auditory aphasic" (Hardy and Pauls, 1959; Wepman, 1960; Monsees, 1961).

Myklebust (1954) hypothesizes that the inconsistent responses of so-called asphasic children may be due to fluctuations in attention, not to differences in sensitivity in auditory stimuli of different frequency. If this is true, then response consistency and the ability to integrate auditory stimuli seem to be two important variables in differentiating the child with a peripheral hearing loss from the child with an auditory aphasia. There is need to develop specific operationally defined procedures for measuring these variables and obtaining clinical information.

There is need to conduct studies on neurological abnormalities which may help differentiate children suffering from peripheral deafness, central deafness, and aphasia. A study of 183 aphasic and deaf children was conducted by Goldstein, Landau, and Kleffner (1960). They found that 32 percent of the aphasic population had no neurological abnormalities. Approximately 40 percent of both the aphasic and deaf children had abnormal encephalograms. Only gross neurological abnormalities and a few etiological categories were found to differentiate deaf from aphasic children. Goldstein et al. (1960), also reported that an autopsy of a 10-year-old child with aphasia and near-normal hearing revealed extensive bilateral degeneration of the auditory areas of the temporal lobes and of the medial geniculate nuclei. The authors state, however, that a large portion of their aphasic population failed to show clinical or laboratory evidence of extensive lesions. Further research is needed to develop diagnostic techniques for establishing the presence of neurological correlates.

Both central and peripheral disorders can have a disrupting effect on language acquisition. For this reason, it is necessary to more clearly define those behaviors which are associated with central and peripheral auditory disorders, and develop procedures to differentiate between the two conditions.

One approach to differential diagnosis is to place the child in a teaching situation, observe his performance and maintain records of significant behaviors (Monsees, 1961; Bangs, 1961; Reed, 1961). Longitudinal studies of children for whom the etiology of a certain condition is known will add easily verifiable data to the present mass of speculation. There is need to record the behavioral responses which are obtained through diagnostic teaching. Information of this kind may help identify the behaviors and learning characteristics of children with different etiological conditions.

The literature concerning the evaluation of auditory capacity and behavior is quite extensive (Ewing and Ewing, 1944; Hardy and Bordley, 1951; Barr, 1955; Lowell, Rushford, Hoversten, and Stoner, 1956; Reichstein and Rosenstein, 1964). At present, however, clinical observation of behavioral symptoms seems to provide the main basis for assessment, evaluation and diagnosis. There is very little experimental evidence to support these clinical observations, and there is need to conduct systematic investigation in these areas.

To better evaluate children with auditory processing disorders, there is need to develop more effective procedures for presenting auditory stimuli, eliciting responses, and increasing the number of response modes. Greater efficiency in selecting the stimulus mode of input, the method of indicating response, and motivating the child to respond may help reduce the amount of response inconsistency. Recent advances in electronics should help accelerate study of the reception and processing of auditory stimuli.

A thorough assessment and diagnosis often requires skill and training beyond that of the individual practitioner. The otolaryngologist, pediatrician, neurologist, psychiatrist, psychologist, audiologist, speech pathologist, and educator all have specific contributions. Research is needed, however, to develop and recommend administrative alternatives for mobilizing these individuals and creating administrative structures which will permit them to work as a team.

The purpose of auditory training is to help the child make active use of his hearing. This concept has been implemented in working with the residual hearing of hard-of-hearing children but comparatively few studies have been reported concerning the training of auditory perceptual disorders. As in many other areas of perception the literature is heavily weighted in favor of diagnosis and the development of diagnostic procedures. There are a few studies which indicated that some degree of amelioration is possible. Unfortunately, these studies often fail to provide a detailed description of the remedial procedures which are used or the nature of the disorders to which remediation was ap-

plied. The methods section of reported studies often consists of abbreviated lists including such topics as hearing and distinguishing sounds; listening games; listening to contrasting sounds, loud and soft, fast and slow, high and low; following directions; hearing through poetry; listening through stories; music to develop sound discrimination; reproduction of auditory stimuli; and auditory memory training. Despite the lack of detail in reporting remedial approaches, there seems to be clinical agreement that training should be attempted in the deficit areas.

There is need to identify and distinguish the different auditory activities and develop specific remedial approaches for these activities. Table 2, for example, attempting to make distinctions between seven different kinds of auditory activities, provides some direction for developing specific remedial procedures.

A basic research question which needs to be answered is to what extent can remedial effort establish or restore behavioral processes which have been disturbed by organic damage? Luria (1966) has observed that the elementary physiological functions such as sight, hearing, touch, or simple movements are disturbed after lesions of the corresponding areas of the human cerebral cortex and are practically incapable of regeneration. In contrast, however, the more complex forms of mental activity which are affected by local brain lesions are more responsive to retraining. Studies have been conducted to show the success of active attempts of individuals to overcome a defect, and while destruction of certain areas is irreversible, in many cases the complex behavioral processes which have been disturbed by damage may be restored. The way in which this takes place, however, is not clear (Goldstein, 1942; Luria, 1948).

Other practical research questions which need to be answered are: How can the principles of learning be applied to help expedite the remediation of auditory processing disorders? How can we help children attend to auditory stimuli; associate sound with experience; reinforce learning through concrete objects, pictures, gestures, and pantomime; retain sound sequences; reorganize and recall word names; analyze sound sequences; synthesize isolated sounds; retain melody and rhythm patterns; differentiate significant from insignificant stimuli; localize sound; and discriminate between sounds?

REFERENCES

BABENKOVA, S. V. *Special features of the interaction between signal systems in the process of restoration of speech in various forms of aphasia. Proceedings of the Seventh Session of the Institute of Neurology* (USSR, Acad. Med. Sci.; Moscow), 1954.

BABKIN, B. P. Characteristics of the acoustic analyzer in dogs. *Trans. Obsh. Russk. Vrach.* v SPb: 77, 1910.

BANGS, T. E. Evaluating children with language delay. *Journal of Speech and Hearing Disorders,* 1961, *26,* 6–18.

BARR, B. Pure tone audiometry for preschool children. *Acta Otolaryngologica,* 1955, 1–84 (Suppl. 121).

BASTIAN, H. C. *Aphasia and other speech defects.* London: H. K. Lewis, 1898.

BERRY, M. F., and EISENSON, J. *Speech disorders: Principles and practices of therapy.* New York: Appleton-Century-Crofts, 1956.

BUTLER, R. A., DIAMOND, I. T., and NEFF, W. D. Role of the auditory cortex in discrimination of changes in frequency. *Journal of Neurophysiology,* 1957, *20.*

CAMPBELL, A. W. *Histological studies on the localization of cerebral function.* Cambridge, England: Cambridge University Press, 1905.

COSTA, D., MANDEL, I. J., and RAPIN, I. Visual and auditory reaction time under reinforcement conditions in children at a school for the deaf. *Asha,* 1962, *4,* 415.

DALE, D. M. C. *Applied audiology for children.* Springfield, Ill.: Charles C Thomas, 1962.

DICARLO, L. M. Differential diagnosis of congenital aphasia. *Volta Review,* 1960, *62,* 361–364.

DICARLO, L. M., and BRADLEY, W. H. A simplified auditory test for infants and young children. *Laryngoscope,* 1961, *71,* 628–646.

DIX, M. R., and HALLPIKE, C. S. The peep show. *British Medical Journal,* 1947, *2,* 719–723.

DOUGLASS, F. M., FOWLER, E. P., Jr., and RYAN, G. M. *A differential study of communication disorders.* New York: Columbia-Presbyterian Medical Center, 1961.

DOWNS, M. P. *The auditory development of the normal child and its relation to testing and therapy for the hearing handicapped child.* Tel Aviv: Micha Society, 1960.

DRAKE, C. *P.E.R.C. auditory discrimination test.* Sherborn, Mass.: Perc. Ed. and Research Center, 1965.

EWING, I. R., and EWING, A. W. B. The ascertainment of deafness in infancy and early childhood. *Journal of Laryngology and Otolaryngology,* 1944, 59, 309–333.

FERRIER, D. *The functions of the brain.* London: Smith, Elder & Co., 1876.

GOLDBERG, I. M., DIAMOND, I. T., and NEFF, W. D. Auditory discrimination after ablation of temporal and insular cortex in cats. *Fed. Proc.* 16, 1957.

GOLDSTEIN, K. *Aftereffects of brain injuries in war.* New York: Grune & Stratton, 1942.

GOLDSTEIN, K. *Language and language disorders.* New York: Grune & Stratton, 1948.

GOLDSTEIN, R., LANDAU, W. M., and KLEFFNER, F. R. Neurologic observations in a population of deaf and aphasic children. *Annals of Otology, Rhinology, and Laryngology,* 1960, 69, 756–767.

GRANIT, R. *Receptors and sensory perception.* New Haven: Yale University Press, 1955.

GUILFORD, F. R., and HAUG, C. O. Diagnosis of deafness in the very young child. *Archives of Otolaryngology,* 1952, 55, 101–106.

HARDY, W. G., and BORDLEY, J. E. Special techniques in testing the hearing of children. *Journal of Speech and Hearing Disorders,* 1951, *16,* 123–131.

HARDY, W. G., and PAULS, M. D. Significance of problems of conditioning in G.S.R. audiometry. *Journal of Speech and Hearing Disorders,* 1959, *24,* 123–126.

HILGARD, E. R. *Theories of learning.* (2d ed.) New York: Appleton-Century-Crofts, Inc., 1956.

HIRSH, I. J. The measurement of hearing. New York: McGraw-Hill, 1952.

KEASTER, J. A quantitative method of testing the hearing of young children. *Journal of Speech Disorders,* 1947, 12, 159–160.

KORST, L. O., and FANTALOVA, V. L. Characteristics of disorders of certain cortical functions in patients with tumors of the temporal and occipital lobes of the brain. *Problems in Modern Neurosurgery,* Vol. 3. Moscow, 1959.

KRYZHANOVSKII, I. I. Conditioned reflexes after extirpation of the temporal regions of the cerebral hemisphere in dogs. (Candidate Dissertation), Military Med. Acad. St. Petersburg, 1909.

LOWELL, E. L., RUSHFORD, G., HOVERSTEN, G., and STONER, M. Evaluation of pure tone audiometery with preschool age children. *Journal of Speech and Hearing Disorders,* 1956, *21,* 292–302.

LURIA, A. R. *Higher cortical functions in man.* New York: Basic Books, 1966.

LURIA, A. R. *Restoration of functions after brain injury* (Russian). Moscow: USSR Academy of Pedagogic Sciences Press, 1948. (English translation, London: Pergamon Press, 1964).

MECHAM, M. J. Shaping adequate speech, hearing and language behavior. In M. J. Mecham, M. J. Berko, F. G. Berko, and M. F. Palmer (eds.), *Communication training in childhood brain damage.* Springfield, Ill.: Charles C Thomas, 1966, 115–143.

MECHAM, M. J., BERKO, M. J., BERKO, F. G., and PALMER, M. F. (eds.), *Communication training in childhood brain damage.* Springfield, Ill.: Charles C Thomas, 1966.

MEYERSON, L., and MICHAEL, J. L. *The measurement of sensory thresholds in exceptionl children: An experimental approach to some problems of differential diagnosis and education with special reference to hearing.* Houston, Tex.: University of Houston, 1960.

MILLER, G. A. *Language and communication.* New York: McGraw-Hill, 1951.

MILLER, G. A., HEISE, G. A., and LICHTEN, W. The intelligibility of speech as a function of the context of the test materials. *Journal of Experimental Psychology,* 1951, *41,* 329–335.

MILLER, N. E., and DOLLARD, J. *Social learning and imitation.* New Haven: Yale University Press, 1941.

MONROE, M. *Children who cannot read.* Chicago: University of Chicago Press, 1932.

MONSEES, E. K. Aphasia in children. *Journal of Speech and Hearing Disorders.* 1961, *26,* 83–86.

MYKLEBUST, H. R. Aphasia in children. In L. Travis (ed.), *Handbook of speech pathology.* New York: Appleton-Century-Crofts, 1957.

MYKLEBUST, H. R. *Auditory disorders in children: A manual for differential diagnosis.* New York: Grune & Stratton, 1954.

NIELSEN, J. M. *Agnosia, apraxia, aphasia: Their value in cerebral localization.* (2d ed.) New York: Hafner Publishing Co., 1948.

O'NEILL, J. J. *The hard of hearing.* Englewood Cliffs, N.J.: Prentice Hall, 1964.

O'NEILL, J. J., OYER, H. J., and HILLIS, J. W. Audiometric procedures used with children. *Journal of Speech and Hearing Disorders,* 1961, *26,* 61–66.

PENFIELD, W., and RASMUSSEN, T. *The central cortex of man.* New York: Macmillan, 1950.

PENFIELD, W., and ROBERTS, L. *Speech and brain mechanisms.* Princeton: Princeton University Press, 1959.

PRONOVOST, W., and DUMBLETON, C. A picture-type speech sound discrimination test. *Journal of Speech and Hearing Disorders,* 1953, *18*(3), 260–266.

REED, M. Hearing and speech disorders. *Cerebral Palsy Bulletin,* 1961, *3,* 52–56.

REICHSTEIN, J., and ROSENTEIN, J. Differential diagnosis of auditory deficits: A review of the literature. *Exceptional Children,* 1964, *31*(2), 73–82.

SEMERITSKAYA, F. M. Rhythm and its disturbance in various motor lesions (Candidate Dissertation, Inst. Psychol.). Moscow, 1945.

SIEGENTHALER, B. M., PEARSON, J., and LEZAK, R. J. A speech reception threshold test for children. *Journal of Speech and Hearing Disorders,* 1954, *19,* 360–366.

SILLS, S. An evaluation of two techniques for screening the hearing of preschool children. Unpublished master's thesis, Hunter College, 1962.

SLOANE, H. N., and MACAULAY, B. D. Teaching and the environmental control of verbal behavior. H. Sloane and B. MacAuley (eds.), *Operant procedures in remedial speech and language training.* Boston: Houghton Mifflin Co., 1968.

SOKOLOV, E. N. *Perception and the conditioned reflex.* Moscow: Moscow University Press, 1958.

SOLOMON, S. Evaluation of hearing acuity in very young children. Unpublished master's thesis, Hunter College, 1962.

SORTINI, A. J. Hearing evaluation of brain-damaged children. *Volta Review,* 1960, *62,* 536–540.

STRENG, A., FITCH, J., HEDGECOCK, L. D., PHILLIPS, J. W., and CARRELL, J. A. *Hearing therapy for children.* (2d rev. ed.) New York: Grune & Stratton, 1958.

TRAUGOTT, N. N. Sensory alalia and aphasia in childhood. *Research Abstracts on Medicobiological Sciences.* No. 1 (Izd. Akad. Med. Nauk SSSR; Moscow), 1947.

UTLEY, J. *What's its name: A guide to speech and hearing development.* Urbana, Ill.: University of Illinois Press, 1950.

WEPMAN, J. M. Auditory discrimination, speech and reading. *Elementary School Journal,* 1960, *60,* 325–333.

WEPMAN, J. M. *Auditory discrimination test.* Chicago: Language Research Associates, 1958.

WERNICKE, C. *Der aphasische symptom komplex.* Breslau: Cohn and Weigert, 1874.

WHITEHURST, M. W. Testing the hearing of preschool children. *Volta Review,* 1961, *63,* 430–463.

WOLFE, W. G., and MACPHERSON, J. R. *The evaluation and development of techniques for testing the auditory acuity of trainable mentally retarded children.* Austin, Tex.: University of Texas, 1959.

YEDINACK, J. G. A study of linguistic functioning of children with articulation and reading disabilities. *Journal of Genetic Psychology,* 1949, *74.*

CHAPTER 3

VISUAL PROCESSING

This chapter is concerned with central dysfunctions in which the subject can see, but experiences difficulty in: (*a*) visually examining the individual details of an object; (*b*) identifying the dominant visual cues; (*c*) integrating or combining individual visual stimuli into simultaneous groups and obtaining meaning from the object; (*d*) classifying the object in a particular visual category; and (*e*) comparing the resulting visual hypothesis with the actual object as it is perceived. This process for receiving, integrating, and decoding or interpreting visual stimuli has been commonly referred to as visual perception.

For many years, traditional psychology and education viewed visual perception as a passive process, which was mainly dependent upon the stimuli reaching the sense organs. Thus, the visual perception of an object simply resulted in the direct and passive visual reflection of that object. There are a number of research studies, however, which indicate that the processing of visual stimuli is not only a more complex act than previously supposed, but a highly active and investigatory process as well. Table 5 presents an outline for the processing of visual stimuli. These eight stages help point out the complexity and the active nature of visual processing operations. The subject actively scans the object, identifies the significant visual cues, and attempts to integrate them into a simultaneous spatial construct. The visual image is tentatively put into a category and compared with the actual object as it is perceived. If the visual image is consistent with the object, the person terminates the perceptual activity. If the visual image and the actual facts are in disagreement, corrections are introduced into the previous visual hypothesis.

Objects which are unfamiliar to the viewer may require all eight stages. In contrast, the process for perceiving familiar objects is more brief. Complex visual objects which are familiar to the viewer can be identified from one dominant sign, and verification that the image is correct takes place almost instantaneously (Gibson, 1966).

It is important to note that the central processing of visual stimuli actually begins with the identification of visual cues. While operations such as receiving visual stimuli, orienting the head and eyes to the light source, and scanning the object do not represent central processing operations in the strictest sense of the word, they are intimately concerned in the process of perception.

The Mechanism for Processing Visual Stimuli

In reviewing the research, it seems that the visual processing mechanism consists of three major parts: (*a*) The ocular-musculature as an adjustor; (*b*) the eye as a transducer; and (*c*) the cortex as a visual processor. This section will present a brief discussion of these mechanisms and their relationships to the central processing of visual stimuli.

The Ocular-Musculature as an Adjustor

The significance of eye movement in "visual perception" has not been made clear. Both autonomic and voluntary movements are intertwined in the complex act of seeing. Morgan and King (1966) point out that the ocular-musculature is never at rest. According to Yarbus (1956), lasting visual impressions require minute muscular movements of the eyes which continually alter the position of the image on the retina. These frequent eye jumps cause the object to be viewed by new receptors. Thus, the electromagnetic

Table 5.—Steps in the Processing of Visual Stimuli

1: Receive visual stimuli:
2. Orient head and eyes to the light source:
3. Scan the object:
4: Identify the dominant visual cues:
5: Integrate the dominant visual cues:
6: Tentatively classify the object in a visual category:
7: Compare the resulting visual hypothesis with the actual object as it is perceived:
8: Confirm comparison or introduce corrections into the previous visual hypothesis:


Primary source of light produced the initiating
Reflected from some object
electromagnetic signal
refracted through the lens and corneal fluid to the retina
Transduced through rods and cones
chemical signal
Transduced to the optic nerve
electrical signal
Transmitted to the visual cortex
visual signal

Figure 3.—The Transduction of Visible Stimuli

excitation is spread over a fairly wide area of the retina which probably prevents fatigue of the receptor elements. These extremely small tremors of the three pairs of eye muscles (saccadic movements) are essential to vision (Riggs, Ratliff, Cornsweet, and Cornsweet, 1953). When an image falls on a single immobile point of the retina for a period of 2 to 3 seconds, it will no longer be perceived (Yarbus, 1956).

In addition to the constant automatic or reflex movements of the eye muscles, the eye actively searches for information in the visual world by scanning the optic array (approximately 140° horizontally and 120° vertically). Scanning is accomplished by using the three pairs of opposing eye muscles to move the eyeball. An exact and subtle nervous system maintains the center of fixation upon both foveas at the same time. Gibson (1966), in comparing the eyes to the front wheels of an automobile, says that the eyes would move as if connected by an invisible tie rod. Failure to develop control may result in poor information transmission. "Each eye can no longer explore the array on its own, for the saccadic movements have what is called compulsory conjugation. Rotations are equal in angle and synchronized. This is necessary, of course, if the fixations are to coincide at the end of a movement. The same is true of pursuant movements." (Gibson, 1966, p. 177)

The Eye as a Transducer

For visual perception to occur, light waves must be transmitted to the central cortex, which presupposes a certain degree of sensitivity of the retina to visible light. The human eye is responsive to two classes of visible light. The first represents the original or primary source of light, which is transmitted through the atmosphere, and impinges directly upon the eye. The second class of visible light is referred to as luminance or reflected light, which is the light that is reflected by leaves, houses, automobiles, pictures, lines, printed words, and other tangible objects which are not primary sources of light.

Figure 3 is a schematic description of the changes which occur to the original light waves emanating from the light source until they reach the visual cortex as electrical signals. The primary or secondary light source produces electromagnetic signals which are transmitted through the lens and corneal fluid to the retina. The ciliary muscles attached to the ligaments holding the lens in place contract so that the lens becomes flat and thin to bring far objects into focus on the retina. When the ciliary muscles are relaxed, the lens becomes curved and thick and near objects are brought into focus on the retina. The rods and cones in the retina transduce the electromagnetic signals into chemical signals which are in turn transduced into electrical signals at the optic nerve. The electrical signals are then transmitted to the occipital cortex.

The Cortex as a Visual Processor

Information on the structure and function of the central nervous system mechanism for visual stimulus processing comes from two main sources: Direct stimulation of the exposed brain, and the study of patients with brain lesions. Extensive investigations of the results of electrical stimulation of the exposed brain are reported by Penfield (1958), Penfield and Jasper (1954), and Penfield and Roberts (1959). Extensive reviews of the literature on cerebral dominance were compiled by Kirk (1935) and Downer (1962). Most of the data are gathered from adults, and consequently few conclusions

can be drawn about the development of the visual system.

It is generally accepted that the retina is an outward extension of the cerebral cortex. The retina of each eye is divided along a vertical line into two halves. Optic nerve fibers transmit sensations from the retina in each eye to the visual cortex of the brain. The optic nerves which lead from the outside halves of each eye transmit sensations to the homolateral hemisphere of the brain. Nerve fibers from the interior or nasal halves of the retina cross at the optic chiasma and transmit sensations to the contralateral hemisphere. The net result is that visual information is relayed through each eye to both hemispheres, and the electrical impulses arising from the retina are transmitted along the optic fibers to the lateral geniculate body and nuclei of the thalamus to the visual areas (areas 17, 18, and 19) of the occipital region of the cerebral cortex.

Studies of animals have shown that extirpation of the occipital portion of the cortex markedly impairs visual discrimination (Lashley, 1930, 1942; Klüver, 1927, 1937, 1941; Chow, 1952; Mishkin and Pribram, 1954). Complex discrimination tasks which required selective discrimination in size, shape, and color were most seriously disturbed by damage to the occipital lobes, whereas the ability to recognize light was not destroyed. These findings suggest that one of the more important functions of the occipital cortex is the analysis and synthesis of visual stimuli. It should be noted, however, that postoperative disturbances may be dependent not only upon destroyed striate area, but upon other regions of the cortex as well. In studying the discrimination of complex visual patterns in rats, Kirk (1936) found that the habit of reaction to a complex visual pattern in terms of relearning is closely proportional to the extent of the lesion, and is not mainly dependent upon the destruction in the area striata or the posterior cortex.

The frontal lobes have been identified as an anterior ocular-motor center for voluntary eye movement. A lesion in the frontal lobes, which disrupts the voluntary control of the eyes, will interfere with active visual investigatory activity (Holmes, 1919, 1938). There is need, however, to study the interrelationships between the occipital and frontal lobes with respect to the processing of visual stimuli.

Many individuals consider motor development to be an important part of the development of vision. Getman (1965) notes that development of vision is dependent upon the motor development of the child. Flavell (1963) described Piaget's theory of sensory-motor development in relation to the development of vision. Dunsing and Kephart (1965) also emphasize the importance of what they term the perceptual motor match.

These is need to conduct additional research on the mechanism for processing visual stimuli. The functioning of the ocular-musculature as an adjustor and the cerebral cortex as a central processor need further investigation. Also, it is important to clarify the interrelationships between the two.

Visual Processing Tasks

Because of the vague and ambiguous terminology and the sketchy procedural descriptions in many research reports, it is difficult to identify specific visual processing tasks. Furthermore, a false distinction seems to have been made between peripheral acuity and central processing perceptual tasks. In reviewing the literature it becomes obvious that there is no clear-cut boundary between visual acuity and the processing of visual stimuli. Instead, there is a continuum of tasks. Nevertheless, for purposes of reporting the research literature, this continuum has been divided into two main classes.

The first class of visual processing tasks consists of the ocular-motor tasks, which are responsible for the reception of visual stimuli. The second class of visual processing tasks are cognitive tasks, which require the analysis and synthesis of visual information.

The discussion of each task will include descriptions of the: (*a*) procedures used to assess performance on these tasks; (*b*) different kinds of visual processing dysfunctions; and (*c*) training procedures which have been developed to ameliorate or compensate for these dysfunctions.

Ocular-Motor Tasks

Ocular-motor tasks include distinguishing light from no light, seeing fine detail, binocular fusion, convergence, and scanning. The relationships between these ocular-motor tasks and visual processing have not been fully explored. According to Gibson (1966), the eye receives visible light and transmits visual information. There is evidence that performance in ocular-motor tasks affects visual processing. As previously noted, the retina is an extension of the cerebral cortex. The importance of the autonomic saccadic eye movements to the continual registration of retinal images has also been discussed. This section will describe the major monocular and binocular motor tasks which have been identified and which are related to the processing of visual stimuli.

861-752 O—70——8

1. *Distinguishing Light From No-Light.* Sensitivity to light is a prerequisite to "visual perception." Without this sensitivity electromagnetic signals cannot be transduced to chemical and electrical signals which are directed to the occipital cortex. The usual procedure for determining sensitivity to light is to present a visible stimulus to discover if the subject can detect light from no-light.

2. *Seeing Fine Detail.* The amount of visual stimuli which is processed is determined, in part, by the detail which can be discriminated. Tests of visual acuity attempt to assess the extent to which the eye or eyes can discriminate details of a series of objects gradually decreasing in size.

Most tests of visual acuity are presented first to one eye and then to the other. Acuity is typically measured by performance on a standard set of printed designs. The Snellen chart, designed to test vision for objects 20 feet from the person, measures responses to the printed letter E. Children are asked to tell which way the "fingers" of the E are pointing, or to point with their own finger or hand in the same direction. The apparent size of an object varies with its distance from the eye. By varying the actual size of the object, symbol, or picture, the distance at which the person can effectively see and report details can be estimated. The Landolt ring, a form shaped like a capital C, and the parallel bars, two rectangles, are also used to assess visual acuity. The person being tested indicates the direction in which the white space points in the Landolt ring, or whether he sees two bars or one. Unless the child has shown such behavior as squinting, holding things close to his eyes, and not recognizing objects and people across a room, examinations for visual acuity are typically not performed until a child enters school.

Organic conditions such as cataracts, tumors, glaucoma, lack of color vision, or damage to the retina will reduce sensitivity to visual stimuli. Failure to respond to light can also be due to damage to the occipital cortex. Total destruction of the occipital cortex of both hemispheres has been found to result in central blindness.

Partial lesions may lead to a blind spot or visual loss in part of the visual field. Teuber (1960a) has shown that the visual area which remains intact continues to function and helps compensate for the deficit caused by the constricted visual field. Luria (1966a) cites case studies by Potzl where lesions of the projection divisions of the occipital cortex are accompanied by hemianopsia, in which the visual field became blurred for a short time or was partially lost, and compensatory eye movements tended to occur.

When damage occurs to the occipital cortex in only one hemisphere, partial blindness, hemianopsia, will occur in both eyes. Half the visual field in both eyes will be affected (Morgan and King, 1966). If the right occipital cortex is damaged, for example, blindness will occur in the outside half of the left visual field and in the nasal half of the right visual field. A lesion which extends outside the primary visual cortical fields, however, may produce different effects. A unilateral lesion involving the parieto-occipital area may result in a hemianopsia which is not compensated for by eye movements. To test the visual fields, the subject is asked to gaze at a fixed point and report the appearance of a second object at the periphery (Luria and Skorodumova, 1950). These conditions are medical problems and should be diagnosed and treated by an ophthalmologist and neurologist.

3. *Binocular Fusion.* Since the visual fields overlap, visual information coming to the brain from the overlapping portions of the visual fields must be integrated into a single set of visual information. The process of integrating the visual fields is binocular fusion.

If one eye is different in refractory power, the quality of the images reflected upon the retinas will differ, and so will the information transmitted to the cortices. This can result in interference between the two sets of information. Some children apparently suppress the image coming from the less effective eye. After several years, information coming from that eye may not be interpreted, and the end result is that the eye is not functioning. If a structural muscle defect prevents an eye from rotating sufficiently, surgery by an ophthalmologist may be needed.

Screening tests for binocular fusion include the Keystone Telebinocular, the Ortho-Rater, and the Massachusetts Vision Test. Care must be taken to ensure that the child knows what is expected of him during the test. Poor binocular fusion is a medical problem and should be diagnosed and treated by the medical profession.

4. *Convergence.* Muscular imbalance can contribute to poor convergence. Coordinated movements of the eyes are necessary for focusing an image on the fovea. In some cases vision training has been instituted by optometrists, more than any other professional group, in an attempt to provide eye muscle training for the separate and coordinated action of the ocular-motor musculature. Vision training usually consists of eye exercises designed to develop: (*a*) More efficient patterns of ocular behavior for reading and other

ocular tasks, (*b*) divergence-convergence relationships, and (*c*) accommodative facility. Many ophthalmologists do not believe these exercises to be beneficial for most patients.

Taylor and Solon (1957) found that no more than 35 hours of training were needed to develop adequate convergence and about 15 hours to establish comfortable vision, provided that the problem has not been complicated by squinting. Strabismus (crossed-eyes) may require from 75 to 200 hours of training before eyes can be coordinated well enough to respond to a visual examination. Waldstreicher (1962) and Getman and Hendrickson (1966) have also reported remedial programs stressing vision training.

Several methods of vision training have been summarized by Lambeth (1966). For the most part, methods used in vision training are not described in detail and there is little research evidence as to the effectiveness of these training procedures either in ameliorating eye-muscle disorders or improving the processing of visual stimuli.

5. *Scanning.* Gibson (1966) identifies three kinds of scanning tasks in which the eye scans the surface and briefly fixates the image upon the fovea. The first is the natural zig-zag scanning that occurs when a child looks around the room. The second is the pursuit or tracking of a moving object which can be done by one eye at a time. Third is the learned systematic scanning, which is required for reading.

An assessment of scanning performance may include an investigation of both the reflexive and psychomotor systems (Luria, 1966a). The elementary reflex system refers to "* * * the reflex fixation of the point to be perceived by a movement of the eye so that the projection of this point lies in the central visual field (fovea)." (p. 358)

Damage to the brain stem or to the posterior ocular-motor centers of the cortex disturbs the elementary reflex fixation of the point in the central visual field. Moving a light source in an arc at a fixed distance is a way to check if the subject can follow the light by moving his eyes and/or by turning and orienting his head toward the object.

Complex or psychomotor movements refer to the direction of the gaze in response to a verbal instruction or in accordance with the patient's own intention. This is a more complex act than the reflex and requires the use of the second signal system. The second signal system refers to the control of motor behavior through verbal mediators. Luria and Homskaya (1962) state that lesions of the anterior ocular-motor centers or of the frontal lobe lying anterior to these centers may disrupt the eye movements used in the active investigation of an object.

Psychomotor functioning can be measured by asking the patient to look right or left or to the side opposite the object. If the eye movements are the same with respect to speed, range, and steadiness, both the reflexive and psychomotor systems are intact. If the reflex movements are intact, but the psychomotor movements are impaired, the cortical apparatus may be defective. In these cases, special laboratory methods may be used to obtain graphic recordings of eye movements through motion pictures. Instruments may be used to record the movements of the pupil, or to trace the movements of a spot of light reflected from a mirror attached to the cornea.

In testing scanning, the tasks which are presented to one eye at a time may be presented to both eyes at the same time. In an ordinary scanning task, the subject is allowed to visually explore an object or a picture, while his eyes and head movements are observed. Disturbances of the visual field may result in altered search patterns.

A number of studies suggest that frontal lobe damage interferes with the ability to actively search, scan, or examine objects. Lesions in the frontal lobes may result in a "pathological inertia" of the sensory process which apparently interferes with the motor scanning aspects of perception.

Another syndrome, involving frontal lobe lesions, consists of difficulty in examining the picture or object due to passive looking, or slight shifts of the gaze from one point to the next. Little effort is made to actively seek out the identifying signs. When conclusions are reached about the picture, statements are made with confidence, and with no attempts at correction, even if the examiner asks the subject to study the picture more carefully. In some cases detail which is reported from one picture is incorrectly reported in subsequent pictures in which that detail is either absent, or no longer significant.

Apparently, frontal lobe lesions may also result in difficulty in perceiving fast moving objects (Cohen, 1959; Teuber, 1960b). There are four main variables in tasks involving the tracking of a moving object: (1) The amount of head and/or body movement permitted or required; (2) the speed of movement of the target; (3) the angle through which the target is moved; and (4) the response required. When performance on pursuit tasks is being tested, the subject may be requested to hold his head still, or he may be allowed or requested to move his head. The target,

usually a point of light or a small object, is moved through a wide arc in several planes: up-down, side-to-side, closer and farther away. The speed and the angle of the motion may be varied. The subject may be required to name certain details on the object, or merely to follow the target.

It should be noted that most ocular-motor visual processing tasks are closely related to school oriented tasks such as reading, writing, and arithmetic. Referral for examination, therefore, is typically made after failure on some school-related task. The initial observation of a deviant behavior, however, is often made by the parent, a pediatrician, or a teacher or school nurse. Parents notice such aspects as clumsiness, crossed eyes, or poor performance in ball-playing activities. Pediatricians notice physical abnormalities such as crossed eyes and inflammations. Teachers notice behaviors which are typically found on checklists such as excessive blinking, rubbing the eyes, unusual head positions while reading, and physical symptoms, such as tearing, inflammation, and redness of the eyes (Betts, 1954; Bond and Tinker, 1967). There is need to develop checklists for parents and teachers which will help in the early identification of visual problems.

If problems can be identified, then the child can be referred for further assessment and treatment to medical and surgical specialists such as an ophthalmologist and pediatrician or to an optometrist who is trained to correct refractive errors and to detect certain structural and functional abnormalities.

The course of development in atypical individuals has been studied, as has the functioning of individuals who have developed normally and then through injury do not perform adequately.

In order to study the possible effects of brain damage on visual acuity, eye muscle control, and other aspects of visual stimulus processing, a number of investigations have been conducted on cerebral palsied children. Cruickshank, Bice, and Wallen (1957) found that sensory (acuity) defects among cerebral palsied subjects differed little from sensory defects found among the normal population. Performances in visual perception and figure-ground differentiation were slightly poorer among the cerebral palsied, particularly the spastic cerebral palsied. These findings are consistent with the literature reported prior to 1957.

A study of cerebral palsied children by Abercrombie (1960) found that visual processing difficulty with certain tasks may be due to uncoordinated eye movements, or eye-hand incoordination. Faulty eye movements may result in disordered pattern discrimination, which might influence perceptual and visual-motor ability. Abercrombie raised the possibility that developmental lags or arrested development might contribute to problems in these areas.

An investigation of visual processing disorders in normal, spastic, and athetoid children by Wedell (1960) found that impairment is generally limited to the spastic group, where one would expect to find greater impairment in the control of the eye muscles. It is difficult to interpret data from investigations of persons who do not have adequate voluntary control over the skeletal muscular system. Frequently, the type of response required is not explicitly reported in the literature, and consequently statements are made concerning the effect of poor eye-hand coordination on visual perception.

Cognitive Tasks

According to Webster (1965) cognition includes both awareness and judgment. The processing of visual stimuli at the higher cortical levels involves: (a) visual analysis, the separation of the whole into its component parts; (b) visual integration, the coordination of mental processes; and (c) visual synthesis, the incorporation or combination of elements into a recognizable whole. A review of the literature reveals a variety of cognitive tasks requiring the analysis, integration, and synthesis of visual information. While it is recognized that these cognitive tasks are all interrelated, for purposes of presentation, these tasks will be discussed under three major groupings, spatial relationships, visual discrimination, and object recognition.

1. *Spatial Relationships.* Body orientation and spatial relationships are among the first visual processing tasks which the infant and young child begin to acquire, and they are among the last to be fully developed (Piaget, 1935). Spatial awareness includes awareness of space which is located left and right, before and behind, above and below the child's own body. Initially, spatial orientation is ego-centered. Physically, the universe centers around the child. The child gradually develops an awareness of space through input and feedback of the visual, muscular, and vestibular mechanisms.

Dysfunctions in spatial orientation are said to be characterized by: (*a*) Difficulty in left-right discrimination; (*b*) avoidance of crossing the midline of the body with the hand; (*c*) poor depth perception; (*d*) reversals such as, b/d; (*e*) rotations such as, p/d; and (*f*) difficulty in perceiving one's own body in space (Nielsen, 1962; Kephart, 1960).

Children with these dysfunctions typically have trouble placing their hands in a particular position;

matching shapes of geometric figures; maintaining their sense of direction; differentiating vertical from horizontal; copying geometric figures, letters, or numbers; dressing themselves; reading; telling time; or using maps. Situations or tasks which require the differentiation of symmetrically opposite points (b/d) are particularly difficult (Holmes, 1919; Gerstmann, 1924; Head, 1926; Bender and Teuber, 1947, 1948; Critchley, 1953; Hécaen and Ajuriaguerra, 1956).

Spatial disorientation has been attributed to damage to the occipital cortex and to cortical lesions in the infero-parietal and parietal-occipital areas (Nielsen, 1962; Luria, 1966a).

Diagnostic procedures for disorders of this kind include investigation of the motor functions of the hand, the perception of pictures which are disoriented to the reader (upside down, etc.), graphic tests for the spatial arrangement of lines, and correct orientation to the parts of figures forming mirror images. Luria (1966a) reports the development of tests for spatial orientation, utilizing maps, floor plans, or routes by Kolodnaya (1949, 1954) and Shemyakin (1940, 1954). Money (1962) has developed a Road Map Test.

Tests for visual orientation with respect to spatial relationships include tasks such as the construction of patterns or figures from wooden blocks which require visual-spatial preservation and the breaking up of homogeneous parts of a pattern into their component spatial elements (or vice versa).

Tests which measure the complex performance of intellectual operations in space are often focused on the solution of mechanical problems, the ability to trace spatially presented movements, and construction of squares of a certain color using small cubes which have sides painted different colors. (For example, the Block Design Test of the WISC.) Luria suggests that such tests are valuable because they reveal specific types of difficulty and provide qualitative insights into the nature of the child's performance.

The status of development of assessing spatial orientation tasks is primitive. This may be due to the lack of: (*a*) specific descriptions of the significant behaviors which characterize correct and incorrect performance on these tasks; (*b*) information as to the specific stimuli which are effective in eliciting these behaviors; and (*c*) systematic instructions which ensure that the subject understands what is expected of him. There is need for future research to develop behavioral tests, and, if possible, link performance with etiological factors.

Luria (1963) describes a type of "afferent apraxia" in which the subject is able to execute purposive behavior and preserve basic spatial coordinates and kinesthetic sensitivity, but is unable to perform a gesture or action of a symbolic or descriptive nature. Kirk (1961, 1968) has termed this type of communication "motor encoding" or "manual expression." While he can functionally dial a telephone, he may be unable to imitate the dialing of a telephone.

Approaches to compensating for these disorders consist of analyzing the motor act and providing the subject with various aids such as logical explanations of motor sequences which are involved in the movement, utilizing kinesthetic feedback and concrete guides. This approach usually consists of providing general principles in a problem area which can be transferred to other tasks which are somewhat related. Unfortunately, subjects with low verbal ability or who have difficulty generalizing or applying principles may not benefit from this approach as much as subjects who have these skills.

Training activities in spatial disorientation usually begin with teaching the subject to orient to different parts of his body and then to his environment. Strategies for handling spatial problems are introduced whenever possible.

Kephart (1960) writes:

> The early motor or muscular responses of the child, which are the earliest behavioral responses of the human organism, represent the beginning of a long process of development and learning * * *. To a large extent, so-called higher forms of behavior develop out of and have their roots in motor learning. (p. 35)

Kephart (1960) believes that perceptual skills and motor skills should not be considered as two separate activities. Since perceptual skills provide continuous feedback for coordinating motor movements, he believes that perceptual-motor ability should be considered a combined activity.

There are several assumptions which have been made about the interrelationship between visual perception and motor ability. The first assumption is that visual perception is dependent upon learning gross motor skills. This implies that disorders in gross motor skills should be corrected before training in visual perception is undertaken. Another assumption which has been made with respect to the development of perceptual-motor skills is that if a stage of the developmental sequence is not attained, failure will be experienced at the higher stages. Brain-injured children who have failed to develop motor abilities, for example, are believed to have gaps in their developmental patterns.

There is little empirical evidence, however, supporting the hypothesis that basic perceptual-motor training leads to improvement in perceptual-motor abilities or to better academic performance.

There has been an interest in sensory-motor training for many years. As early as 1846, Seguin developed an educational approach which was based on physiological and neurological hypotheses. Seguin divided the nervous system into two parts: the peripheral and the central nervous systems. Disorders in each of these systems would result in isolating the individual and "locking his learning processes."

Seguin attempted to ameliorate these deficiencies through muscular and sensory training. Deficiencies in the peripheral nervous system were trained through sensitivity of the receptors. If the problem was assumed to be in the afferent pathways, the subject was given a series of quickness exercises based upon imitation. If the problem was central, Seguin tried to develop sensitivity by stimulating the cortex into activity with contrasting stimuli. He hypothesized that the stimuli were more apt to be received if they were presented successively at fast rates.

Basic motor skills were taught which began with gross muscle movements and progressed to fine muscle movement. Immobility was taught first, followed by balancing, which was thought to be the primary pivot skill for other movements. Seguin placed great importance on the tactile function, and trained activities such as seizing, holding, letting go, and handling objects appropriately. He also attempted to train taste and smell.

Seguin taught passive, receptive vision of general impressions, and active meaningful perception of seen events. He began by training attention to the stimuli. He used cards, balls, ribbons, fruit, and other objects to train appreciation of color. Awareness of distance, form, and spatial planes was also taught. Listening skills included the passive reception of auditory sounds into meaningful perception of selected sounds and avoidance of other sound impressions. Seguin also provided training in speech, reading, and writing. These activities were primarily done through imitation techniques aided by the sense of touch and the use of flashcards and concrete objects.

More recently, Kephart (1960) has developed a training program for developing perceptual-motor abilities in children who have "suffered breakdowns in perceptual-motor development at one of the earlier stages." Kephart's method is concerned with sensory-motor abilities which are basic to visual-perceptual abilities. Training procedures are based upon specific diagnosis of perceptual-motor development, and are directed at the most basic area of weakness. As performance improves, instruction is shifted to "higher" activities. Kephart emphasizes the importance of developing generalizations, especially motor generalizations, as soon as possible. In order to do this, he has designed activities which appear to transfer to other situations. Although Kephart's program is concerned primarily with perceptual-motor development, it includes verbal language activities of comprehension and expression. For example, he sometimes requests that the child state what he is going to do, tell what he is doing, and afterwards, tell what he has done. Kephart repeats activities which have been mastered in order to reinforce and integrate these activities with new ones.

The sensory-motor training program trains gross motor skill and involves such activities as bounce-to-rhythm, whole body movement, balance-on-board, and movement of differentiated body parts. Chalkboard training improves the development of eye-hand coordination and directionality through scribbling, pursuit, drawing, and copying tasks. Ocular control is developed through ocular pursuit activities, binocular and monocular training, and games which involve a visual component. These activities are designed to establish muscular control of eye movements and to coordinate these movements with other body movements. Form perception training is intended to develop discrimination between object shapes and figure-ground relationships by putting puzzles together, making stick figures, and forming designs. Kephart focuses attention on what the child is doing and the process which he is using. He describes each task in terms of the terminal behavior, the operations necessary to accomplish it, and the rationale of the activity.

It is interesting to note the similarities between the approaches taken by Seguin and by Kephart. Both are concerned with the problem of ameliorating disabilities in the perceptual-motor processes, both approaches are based in part on neurological and psychological hypotheses, and both stress importance of differential diagnosis and the principles of child development. Many of the activities and equipment are similar, such as the springboard—trampoline; nails in a board—pegboard; footprints—stepping stones; and candle—penlight.

The major difference between these two approaches is that Seguin directed his method toward severely mentally deficient children, whereas Kephart has developed his program for children with less severe problems. Because of the severity of the cases with which he

worked, Seguin placed less emphasis on the verbal aspect of the program than did Kephart.

Painter (1966) studied the effect of a rhythmic and sensory motor-activity program on the perceptual-motor spatial abilities of kindergarten children. The purpose of this study was to investigate the effects of the program on body image, perceptual-motor integration, and the psycholinguistic competence of kindergarten children. Twenty-one half-hour training sessions were given to the experimental group extending over a period of 7 weeks. The children were seen three times a week. The program was carefully sequenced using theoretical constructs suggested by Barsch (1963) and Kephart (1960). Thirty-eight activities were related to nine of 12 movement areas of Barsch's Movegenic Theory. The nine included: Visual dynamics; auditory dynamics; dynamic balance; spatial awareness; tactual dynamics; body awareness; rhythm; flexibility; and unilateral and bilateral movements. These activities involved: See and move; hear and move; balancing for both sides of the body; awareness of one's body in space; feeling and moving; being aware of, identifying and localizing body parts; movement to auditory rhythmic patterns, change in tempo and movement patterns; and moving one side or two sides of the body.

In addition to the Barsch theoretical constructs, some of Kephart's procedures were used. These included: generalization of rhythmic patterns; the sequencing of unilateral, bilateral, and cross-lateral movement; and the changing of uncoordinated or jerky movements to large, sweeping movements, in which the entire musculature of the body was used. The activities were sequenced so that they would progress from the very simple to the more complex as the children developed skills.

The results of the study demonstrated that a systematic program of rhythmic and sensory activity will: (1) affect the level of ability to draw a human figure; (2) ameliorate the apparent distortion of body image concept; (3) improve visual-motor integrity; (4) improve sensory motor spatial performance skills; (5) improve psycholinguistic abilities; and (6) improve the ability to express ideas motorically. Because a small number of subjects were utilized over a short period of time, Painter considers this investigation as a pilot study. The significant aspect of this study is that a carefully designed program was developed and applied under controlled conditions to bring about significant gains in specific learning and skills, such as body image, perceptual-motor integration; and psycholinguistic competence. The implications of the study may be generalized to preschool children or to children who present problems created by minimal brain dysfunction.

The Frostig Program for the Development of Visual perception (Frostig and Horne, 1964) attempts to develop proficiency in visual perceptual abilities. The authors state that the program is designed to be corrective and preventative. Remediation is based on each of the five areas measured by the Frostig Developmental Test of Visual Perception (1963), i.e., position in space, spatial relationships, perceptual constancy, visual-motor coordination, and eye-hand coordination. The program consists of work sheets and exercises designed to ameliorate any dysfunction in visual perception. The exercises include games and physical activities designed to facilitate visual perceptual development. Physical activities include: (*a*) awareness of body parts; (*b*) right-left differentiation; (*c*) eye-movement exercises; and (*d*) gross and fine motor activities. The five areas of visual perception are trained concurrently.

The efficacy of the Frostig remedial program has yet to be evaluated in its entirety. Rosen (1966) investigated the effect of the worksheets in the remedial program. He compared groups of first grade children. Subjects were tested on the Metropolitan Readiness Test and the DTVP at the beginning of the school year, and retested on the DTVP later in the academic year. The experimental group received 29 days training on the Frostig material (30 minutes per day). The control group received 15 minutes extra reading time, while the experiemental group had 15 minutes subtracted from their daily reading period. Results showed that additional reading time was more important than perceptual training, that improvement in perceptual skills was not reflected in later reading ability, and that perceptual training showed increased perceptual ability as measured by the CDVP. Additional research is needed before the results of this study can be accepted or rejected. It is likely that individual gains in perceptual ability may have been obscured by other factors in the experiment. It should be noted that teaching reading itself is perceptual training. Those who teach children to discriminate an A from a B may have more transfer to reading that those who teach children to discriminate a square from a triangle.

While certain training procedures have been developed and clinical reports of progress have been made, specific training programs have not been related to the various behavioral syndromes characterizing spatial disorders.

2. *Visual Discrimination.* There are several kinds of visual discrimination tasks. In the first, the subject is typically presented with several discrete stimuli, usually

differing in one specific stimulus element or quality of the object, and is required to respond to that quality. This task will be referred to as an object-quality or prominent-feature discrimination task. The second kind of visual discrimination task is one in which a complex stimulus is presented and the number of stimulus elements is increased. When the number of stimulus elements is increased, usually by the addition of irrelevant elements, the task is called "figure-ground differentiation." In the third type, the number of stimulus elements is decreased by omitting a portion of the visual detail of the object. This task is called "visual closure."

Visual discrimination is involved in tasks in which the child is to fit three-dimensional forms into holes (Stanford-Binet Intelligence Scale, 1960), point to the pattern which is different (Columbia Mental Maturity Scale, 1954), draw a circle around the letter form which is different (reading readiness tests), copy designs (Bender Visual-Motor Gestalt Test, 1938).

It is unfortunate that so few studies of "disorders" have reported the procedural details of the tasks and the behavioral responses of the children in performing these tasks. It is not uncommon to read inferences such as "poor visual perception" based on performance on a single task wherein the required response was to draw or reproduce a picture or a pattern. Further, there is confusion with respect to the names given to these tasks. In some instances, two different names have been given to the same task. In other cases, the same name has been applied to two different tasks.

Visual discrimination tasks are operationally defined by the stimuli which are presented, and the response which is required. Visual stimuli can vary along different presentational dimensions including: (*a*) number of visual stimuli which are presented; (*b*) rate of presentation; (*c*) duration of presentation; (*d*) color-hue; (*e*) brightness; (*f*) contrast; (*g*) size; (*h*) shape; (*i*) sequential order; and (*j*) motion. Various combinations of presentational aspects define different visual discrimination tasks. Visual stimuli may be presented either at the same time (simultaneous), or at different times (successive). Discrimination becomes more difficult as the total number of visible units in the stimuli increases. The discrimination task is made easier if the subject is familiar with the set of significant features, and if both members of the pair are presented at the same time.

A. Selection on the Basis of Dominant Features: Children have been shown to discriminate dominant features (object qualities, attributes) such as: color and form (Corah, 1964); shape (Ling, 1941); pattern (Dornbush and Winnick, 1966); letter-like forms (Gibson, Gibson, Pick, and Osser, 1962); Greek letters (Shepard, 1957); size (Kagan and Lemkin, 1961); position (Lubker, 1964); brightness (Clifford and Calfin, 1958); area (Welch, 1939); and horizontal versus vertical direction of a stripe (Jeeves, reported in Munn, 1965).

The amount of time it takes an individual to perceive visual stimuli is another variable which affects visual discrimination. A discrimination task usually becomes more difficult as the presentation time of a stimulus item decreases. The effects of intervals shorter than one-half second are complex due to the time needed for the human eye to fixate on more than one point to gain sufficient visual information (Gibson, 1966). A child may be able to successfully complete the tests provided he is given time to study and examine the stimulus material. When the stimulus is presented for only a brief time, as in tachistoscope, he may be unable to perform the task.

A dysfunction in perceptual speed is characterized by extended examination of pictures, perception of only one element at a time, and difficulty in selecting and relating the necessary identification signs. It may be accompanied by an ataxia of gaze and uncertain conclusions about what is seen. These symptoms may be due to lesions of the occipital lobes.

The discrimination of objects differing in one dominant feature may be the first step in the complex task of processing visual stimuli. Simple discrimination is also one of the first steps in assessing performance in visual processing tasks in order to determine which aspect of the complex process is disturbed.

Pictures are presented to the subject and he is asked to name them. This approach is useful in identifying subjects who: (*a*) recognize objects or pictures; (*b*) perceive only one cue; (*c*) identify and synthesize the essential cues of a picture; (*d*) relate them to other signs; and (*e*) draw conclusions from the single cue which has been perceived. Teuber (1962) found that certain visual symptoms appear to be more marked after right than left hemispheric involvement. This suggests that many of the visual perceptive skills are mediated in the right hemisphere.

There are few procedures which have attempted to train discrimination activities. Montessori (1964), for example, developed a set of didactic materials for educating the sense of vision. These materials include a set of objects which differ in size, thickness, length, form, and color. The purpose of the exercises using these materials is to train discrimination of objects and forms. Montessori states that by utilizing the tactile

modality, visual recognition of forms is greatly improved.

The application of operant techniques and the use of programmed instruction has been used to ameliorate visual-perceptual disorders. A fading procedure was used by Moore and Goldiamond (1964) to establish visual discrimination in preschool children. Staats and Staats (1963) describe the use of reinforcements to improve discrimination of visual verbal stimuli and reading ability. Programming a task according to operant principles may involve the following steps:

1. Select a form or class of behavior that seems essential to the performance of some everyday (academic) task.
2. Describe the behavior in detailed, objective terms, that is, establish a reliable set of clearly specified criteria.
3. Prepare a sequence of materials and establish a procedure aimed at enabling the individual to perform such a task.
4. Find children who cannot perform as required (even though they possess the necessary biological equipment) or who perform it poorly, and give them training on the sequence.
5. Modify the sequence systematically until the subjects perform according to criterion.

Seguin (1907) attempted to ameliorate visual perceptual deficiencies through muscular and sensory training. Deficiencies in the peripheral nervous system were trained through sensitivity of the receptors. If the problem was assumed to be in the afferent pathways, the subject was given a series of quickness exercises based upon imitation. If the problem was central, Seguin tried to develop sensitivity by stimulating the cortex into activity with contrasting stimuli. He hypothesized that the stimuli were more apt to be received if they were presented successively at fast rates.

There have been a few studies which have attempted to differentiate how the visual-perceptual performance of brain-injured persons is distinguished from the performance of non-brain-injured individuals. Many of these studies, however, attempted to compare brain-injured subjects with familial retardates rather than with normal subjects (McMurray, 1954; Halpin, 1955; Hunt and Patterson, 1958). Total score on a test or error scores are often reported, obscuring qualitative evidence. There is a need for more detailed information which describes the nature of visual-processing performances required in the tasks. Elkind, Koegler, Go, and Van Doorninck (1965) studied the effects of perceptual training on unmatched samples of brain-injured and the familial retarded children. Both groups improved in the perception of ambiguous figures; however, the brain-injured group required more trials and reached a lower level of performance than did the familial retarded group.

B. Figure-Ground Differentiation: The concept of figure-ground differentiation refers to the distinguishing of an object from its general sensory background. According to Rubin (1921) the figure appears to be in front, and the ground consists of those stimuli which are behind the figure. This represents a special type of visual discrimination task. The subject must attempt to locate a stimulus, which has been identified as the figure, despite the distracting effects of competitive visual stimuli which are seen simultaneously by the subject. In order to accomplish this task the subject must attempt to form and retain an image of the figure while he is scanning the total stimulus pattern. Children have been shown to perform this type of task (Munn and Steining, 1931; Gellermann, 1933).

The response modes which are required of subjects in figure-ground tasks vary from task to task. Superimposed figure tasks may require the subject to indicate his choice of figure by drawing around it or on it with a pencil. Some tasks require counting the number of times a certain object occurs. Luria (1966a) describes a task in which the subject is shown pictures which have been scribbled over or superimposed on one another, and is asked to identify the outlines. Luria cites Poppelreuter's (1917, 1918) tests for superimposed figures as examples of early tests for figure-ground discrimination.

The Frostig Developmental Test of Visual Perception (Frostig, 1963) contains a figure-ground task. The remedial program based on the test (Frostig and Horne, 1964) contains especially developed sequentially ordered tasks which presumably train the performance.

More recently, Metzger (1953) has developed stationary optical designs in which searching and scanning for a given figure is made difficult by additional lines.

Some researchers have related poor performance in figure-ground tasks to damage in certain parts of the brain. Strauss and Lehtinen (1947) hypothesized, and Teuber and Weinstein (1956) found that damage to the frontal lobes results in figure-ground confusion, distraction by irrelevant stimuli, and a return to the most prominent element of the stimulus field. Because the figure-ground tasks in the classroom have not been carefully identified and described, little effort has been made to develop training programs to improve performance on a variety of these tasks.

C. Visual Closure: A visual closure task is another special kind of visual discrimination in which the subject is asked to recognize or identify an object, despite the fact that the total visual stimulus is not presented.

The significant features which define and allow recognition of an object or picture are typically given, but not the entire set of features or the entire pattern. He may be required to demonstrate his recognition of it by naming; pointing to incomplete forms of a specified object among a variety of figures; or making the form complete. This task is found in the "Incomplete Man" item on the Stanford-Binet Intelligence Scale, in which the child is asked to complete the picture by drawing. The "Visual-Closure" subtest of the revised Illinois Test of Psycholinguistic Abilities (Kirk, McCarthy, and Kirk, 1968) is another example of a test measuring recognition of incomplete objects. There is need to develop and evaluate training programs to determine whether or not dysfunctions in visual closure can be ameliorated.

3. *Object Recognition (Visual Agnosia).* The behavioral phenomenon called "central blindness, mental blindness, or visual agnosia" was described nearly one hundred years ago. Case studies of animals and humans were reported in which the subjects were no longer able to recognize objects, despite the fact that the elementary functions of seeing objects, describing parts of objects, reproducing objects, and avoiding obstacles remained intact (Jackson, 1876; Munk, 1881; Charcot, 1886, 1887; and Lissauer, 1889). This condition was linked with lesions of the occipital and parietal regions of the brain. More recently, animal studies have supported earlier findings that damage to the occipital cortex disturbs the processing of visual stimuli. While the discrimination of shape and color is impossible, the elementary visual functioning such as the gross recognition of light is not so severely affected (Lashley, 1930, 1942; Kirk, 1936; Chow, 1952; Mishkin, 1954; and Mishkin and Pribram, 1954).

In the history of disturbances of visual processes, disagreement existed as to whether visual imperception represented a complex symbolic dysfunction, or a disturbance in visual memory. Recent research suggests, however, that visual agnosia is a dysfunction of the higher cortical analysis and synthesis of visual images.

While the recognition of objects through the visual channel may be impaired, a child may be able to recognize objects through the sense of touch. In cases where damage to the occipital lobes is less severe, tasks which require the recognition of simple objects are completed more easily than are complex visual tasks which are composed of many visual stimuli. For this reason it is necessary to assess the range of tasks from simple to complex.

The underlying disorder seems to be one in which visual processing is incomplete, because the subject is unable to synthesize visual stimuli and integrate them into a unified whole. Instead, the subject focuses on one dominant visual stimulus and guesses. Attempts to integrate two or more stimuli are usually unsuccessful. Birenbaum (1948) found that it is not uncommon for subjects with language ability to attempt to compensate for visual agnosia by converting their visual analysis into a verbal analysis. Subjects were as effective in interpreting parts of pictures as they were in interpreting complete pictures. Narration was used as the mode of compensation.

In mild cases of occipital and occipito-parietal involvement, few signs of visual agnosia appear. For example, the child may be able to distinguish between objects and pictures. In these cases the child may have difficulty perceiving more than one object at a time, thus affecting the integration of elements into groups. Difficulty in shifting the gaze from one object to another is another symptom of mild involvement (Holmes, 1919; Hécaen and Ajuriaguerra, 1952; Yarbus, 1961).

Pavlov (1949) hypothesizes that failure to deal simultaneously with two or more stimuli may be due to an inhibitory effect that occurs when one stimulus creates a point of excitation which in turn inhibits the excitation of the second stimulus.

A number of behavioral symptoms have been associated with lesions of the frontal lobes. These include: (*a*) inability to inhibit reactions from irrelevant stimuli; (*b*) failure to restrain automatic movement; (*c*) apathetic, dull visual investigatory activity; (*d*) failure to select essential cues; (*e*) an inert and immobile gaze; and (*f*) failure to combine details into a whole. In these cases, the visual field is narrowed, and although objects can be seen in the midline or central field of vision, they cannot be seen on the periphery.

With the exception of being able to distinguish color and shape (Pribram, 1959, 1960; Mishkin, 1957), the behavioral symptoms of occipital and frontal lobe involvement appear to be similar. The major difference between the two is that frontal lobe disorders represent dysfunctions of active investigatory activity, whereas, occipital dysfunctions reflect deficits in the analysis and synthesis of visual stimuli.

Luria (1966a) indicates that frontal lobe damage and true optic agnosia may be distinguished by

"* * * careful observation of the passive character of the perceptual activity (in frontal lobe damage) and of the presence of the syndrome of generalized inertia and lack of critical attitude toward performance * * *." (p. 364) The deficit may appear to be the same inability to recognize an object. However, the etiology of this disability is more likely related to a lack of control over the voluntary scanning movements of the eye. In this case, failure to recognize the figure is not an agnosia but is instead an inability to control impulses regulating the eye movements. Scanning is erratic and incomplete. This "frontal syndrome" seems to represent a disturbance in purpose of behavior. This description closely resembles the stereotype of the hyperactive "Strauss Syndrome" child who exhibits impulsivity and perceptual disorders.

There are a number of tests which are currently being used to assess visual processing operations. The Bender Visual-Motor Gestalt Test (1938) was one of the first tests of visual stimulus processing. The designs were adapted by Lauretta Bender from those used by Wertheimer to demonstrate the principles of Gestalt psychology. The Bender test requires the subject to copy designs from a pattern. Failure to copy a design correctly may be due to a deficit in either visual processing or motor performance. It may be necessary, therefore, to assess the visual (perceptual) component separately from the motor reproduction, especially, if a child appears to have a motor problem. In addition to its use as a test of visual-motor perception, the Bender Gestalt has been used for a variety of other purposes, e.g., to diagnose brain injury, emotional disturbances, and as a predictor of school achievement. Noting that between 1938 and 1964 more than 130 publications have been written concerning the test, Koppitz (1964) has integrated the research on children (CA 5–7) into one volume and presents guidelines for objectively scoring the test.

The Frostig Developmental Test of Visual Perception (1963) assesses five areas which the authors consider to be related to academic skills: Figure-ground relationships, eye-hand coordination, form constancy, position in space, and spatial relationships. Motor skills are necessary for adequate performance in each subtest. The DTVP yields a "perceptual quotient" with norms for children from ages 4 to 7. Frostig and Horne (1964) have developed a remedial program for ameliorating deficits in each area.

Several tests require copying a form, using a pencil. The Graham Kendall Memory-for-Designs Test (1960) requires the reproduction of 15 geometric designs from immediate memory. The Developmental Test of Visual-Motor Integration by Beery and Buktenica (1967) requires the copying of a series of forms, with the model at the top of the page and the space for reproduction at the bottom. The Stanford-Binet Intelligence Scale, Form L–M (Terman and Merrill, 1960) contains several copying items: A circle at "Year III," a square at "Year V," a diamond at "Year VII." The Purdue Perceptual Motor Survey (Roach and Kephart, 1966) also includes several coyping items, such as a series of loops, circles, and scallops.

Several standardized tests have visual components and varying degrees of motor responses. Most reading readiness or school readiness tests include sections on finding similarities and differences in sets of pictures and forms (Gates, 1939; Hildreth and Griffiths, 1949; Murphy and Durrell, 1965).

Some scales purporting to measure cognitive abilities or intelligence have sections involving visual stimuli. The Columbia Mental Maturity Scale (Burgemeister, Blum, and Lorge, 1954) is a series of visual discrimination items. The so-called performance sections of the WISC (Wechsler, 1949) contain subtests requiring the child to put together flat cut-up drawings (object assembly), indicate missing features (picture completion), arrange pictures in a story-sequence (picture arrangement), copy a geometric design (block design), and copy a different arbitrarily designated symbol underneath a series of random numerals, 1 through 9 (coding). An alternate test is the drawing of a line through a printed linear box maze (mazes).

A recent addition to the test repetoire is the Wechsler Preschool and Primary Scale of Intelligence or WPPSI (Wechsler, 1967) which is similar in content to the WISC, with the exception of the coding task which has been changed somewhat to suit the less skilled motor performance of children 4 through 6½. The Animal House subtest requires the child to associate a different color peg with an animal picture and place a peg of the appropriate color in holes marked with animal pictures. The Goodenough Draw-a-Man Test (Goodenough, 1926) requires the child to draw human figures. The Peabody Picture Vocabulary Test (Dunn, 1951) requires the child to listen to a spoken word and indicate, usually by pointing, the one picture of four presented which was named. In contrast, the Picture Vocabulary item on the Standford-Binet requires the child to name each of a series of pictured objects.

Some tests have been developed with a view toward eliminating both the motor production (copying) of a stimulus and the producing of a free response. The colored Progressive Matrices (Raven, 1956) consists

of a series of items, each arranged in matrix form, usually two by two. In each item some progressive feature can be distinguished, and one cell is missing. It is the child's task to indicate which one of four possible cell-fillers offered to him is appropriate. The Illinois Test of Psycholinguistic Abilities, revised edition (Kirk, McCarthy, and Kirk, 1968) contains five subtests with visual stimuli and motor responses. "Visual sequencing" requires the replacing of form chips in a specified sequence. "Visual reception" requires the subject to attend to one stimulus picture, then choose without the model, the one picture of four which is similar in function. "Visual Association" requires the choice of the one in four pictures which is most closely associated with the stimulus picture, which remains in view. "Visual Closure" requires the subject to point to partially hidden figures such as dogs, shoes, and fish. "Manual Expression" requires the use of gestures to illustrate the use of pictured objects, e.g., hammer, comb, and mirror.

There are a number of disorders which interfere with the processing of visual information. The behavioral symptoms of these disorders, for the most part, are varied and represent different kinds of cognitive operations. For this reason, a differential approach to treatment and compensation is indicated.

In cases of central blindness, where vision is completely lost, it is necessary to intervene with teaching approaches which are used for the blind. If there is partial damage to the occipital lobes resulting in hemianopsia, training manual and ocular sensitivity may help compensate for the visual perceptual deficit.

Case histories report wide differences in the quantity and quality of information reported by subjects with visual processing disorders (Gelb and Goldstein, 1920; Goldstein, 1948; Nielsen, 1962; Luria, 1966a). An attempt should be made, therefore, to ascertain, as accurately as possible, whatever it is that the subject reports he sees, i.e., nothing, blurred objects, outlines, or individual elements of an object. Also, any failure to integrate the elements of the object for purposes of recognition, and the degree of confidence or insecurity with which he reports his impressions should be noted. Self reporting has obvious limitations for use with younger children, as well as older children who have limited expressive language. In any event, an attempt should be made to list the visual processing tasks with which the child is successful, as well as those tasks with which he is unsuccessful.

The number of compensatory modes are limited. Improvement in supplementing visual information through tactile and kinesthetic hand and eye movements is one approach. Training the child to study the outline of an object with hands and/or eye movement may help him in distinguishing the elements of the object, and integrating them into a meaningful recognized whole (Fernald, 1943).

If an individual with a visual processing dysfunction attempts to recognize an object by using language as a compensatory mechanism, he may arrive at an incorrect conclusion, because his verbal reasoning is based on incorrect, incomplete, and unintegrated visual information.

There is need to develop more precise methods for assisting the subject to search for and obtain visual information which, for the most part is correct; as well as develop specific strategies to prevent guessing or drawing of hastly conclusions.

Compensatory mechanisms also may be developed for specific tasks such as reading. For example, alphabet letters can be associated with meaning such as telling the child that "the letter S looks like a snake." This takes advantage of general information possessed by the child to help compensate for a disability in visual processing. Another compensatory method is to teach the child to assemble alphabet letters by showing him the letter or number, analyzing it into its parts, and teaching him to reassemble the parts into the whole. This method also takes advantage of the child's ability to learn, remember, and apply principles as well as become familiar with the motor-kinesthetic sequence of disassembling and assembling the elements of the alphabet letters or numbers.

When the visual perceptual field is narrowed as a result of hemianopsia, the number of visual units which can be seen may be increased by teaching the subject to utilize the residual visual fields by employing active systematic scanning procedures. If, on the other hand, the visual field is narrowed because of simultaneous agnosia, the subject can only process one or two visual units at a time. This disorder is extremely disruptive to the act of reading which requires the rapid analysis and synthesis of words, phrases, and sentences.

Shif (1945) studied subjects with simultaneous visual agnosia, who had a narrowed field of reading. Shif found that in addition to two or three clearly perceived elements, the remainder of the visual field was not entirely empty. Peripheral vision included several poorly differentiated signs. In order to teach the subjects to read, Shif wanted to widen the perception of letters and shift the focus of reading to the right, so the right side of the visual field would become a clear and independent center for reading. He did this by: (*a*) introducing certain groups of letters as single

visual units; (*b*) requiring practice on distinguishing the groups of letters from words; and (*c*) using the groups to produce new short words with the same beginning. This procedure enabled the patients to perceive the fixed groups as a simultaneous entity and the patients ceased to analyze the structure of the word. Instead, the patients were able to concentrate on the right half of the word. Reading became possible and was supplemented by perimetry training.

Most of the research on dysfunctions in the processing of visual stimuli has been done with the adult population. Comparatively little work has been done with children. An important issue which needs to be resolved through future research is whether or not certain dysfunctions can be ameliorated through remediation as opposed to providing compensatory mechanisms for adjustment in the subject's areas of strength. There is need for interdisciplinary idiographic ($N=1$) research between educators, pediatricians, neurologists, ophthalmologists, and psychologists to resolve many of these questions.

Directions for Future Research

The traditional distinction between peripheral and central disorders is often confusing and ambiguous. An effort should be made to clarify this distinction in terms of a continuum of interaction among the three components of the visual processing mechanism: The eye, the ocular musculature, and the brain. A better understanding of the tasks involved in visual stimulus processing will hopefully lead to quicker and more successful educational, psychological, medical, and surgical intervention, as well as preventive measures.

A more accurate description of the ocular-motor tasks known to be related to the processing of visual stimuli is needed. There is need to investigate and identify the more subtle ocular-motor tasks which may be involved such as distinguishing light from no-light, seeing fine detail, binocular fusion, convergence, scanning and tracking, developing spatial relationships, discriminating object qualities, differentiating figure from ground, completing visual wholes, and recognizing objects. The relationships of fine and gross ocular motor tasks to cognitive tasks and to the continuum of visual stimulus processing tasks has not been made clear in the present literature.

Cognitive tasks should be thoroughly investigated along broad types of parameters: stimulus, organism, and response. These investigations should be sequentially programmed so as to bring maximum information from a series of studies. Research is needed to develop different procedures for assessment and training in visual investigatory activity and in improving operations in visual analysis and synthesis.

REFERENCES

Abercrombie, M. L. J. Perception and eye movements: Some speculation on disorders in cerebral palsy. *Cerebral Palsy Bulletin*, 1960, *3*, 142–148.

Barsch, R. Project M.O.V.E. as a model for rehabilitation theory. Paper presented at APA Convention, Philadelphia, 1963.

Beery, K. E., and Buktenica, N. *Developmental test of visual motor integration*. Chicago: Follett, 1967.

Bender, L. A visual motor Gestalt test and its clinical use. *Research Monograph*, American Orthopsychiatric Association, 1938, No. 3.

Bender, M. B., and Teuber, H. L. Spatial organization of visual perception following injury to the brain. *AMA Archives of Neurology and Psychiatry*, 1947, *58*.

Bender, M. B., and Teuber, H. L. Spatial organization of visual perception following injury to the brain. *AMA Archives of Neurology and Psychiatry*, 1948, *59*.

Betts, E. *Foundations of reading instruction*. New York: American Book, 1954.

Birenbaum, G. V. Relationship between the conceptual and structural components of perception. In *Investigations on the psychology of perception*. Moscow: Izd. An SSSR, 1948.

Bond, G., and Tinker, M. *Reading difficulties: Their diagnosis and correction*. (2d ed.) New York: *Appleton*-Century-Crofts, 1967.

Burgemeister, B., Blum, L., and Lorge, L. *Columbia mental maturity scale*. Chicago: World Book Co., 1954.

Charcot, P. *Oeuvres complètes*. 9 vols. Paris: Bureau du Progrès Medicoe, 1886–1887.

Chow, K. L. Further studies on selective ablation of associative cortex in relation to visually mediated behavior. *Journal of Comparative and Physiological Psychology*, 1952, *45*.

Clifford, L. T., and Calfin, A. D. Effect of age on the discriminative learning of color and brightness by children. *American Journal of Psychology*, 1958, *71*, 766–767.

Cohen, L. Perception of reversible figures after brain injury. *AMA Archives of Neurology*, 1959, *81*.

Corah, N. L. Color and form in children's perceptual behavior. *Perceptual Motor Skills*, 1964, *18*, 313–316.

Critchley, M. *The parietal lobes*. London: E. J. Arnold, 1953.

Cruickshank, W., Bice, H., and Wallen, N. *Perception and cerebral palsy*. Syracuse: Syracuse University Press, 1957.

Dornbush, R. L., and Winnick, W. A. The relative effectiveness of stereometric and pattern stimuli in discrimination learning in children. *Psychonomic Science*, 1966, *5*, 301–302.

Downer, J. L. Interhemispheric integration in the visual system. In V. B. Mountcastle (ed.), *Interhemispheric relations and cerebral dominance*. Baltimore: Johns Hopkins Press, 1962.

DUNN, L. M. *Peabody picture vocabulary test.* Minneapolis: American Guidance Service, 1959.

DUNSING, J., and KEPHART, N. Motor generalizations in space and time. In J. Hellmuth (ed.), *Learning Disorders,* Vol. 1. Seattle: Special Child Publications, 1965.

ELKIND, D., KOEGLER, R. R., Go, E., and VAN DOORNINCK, W. Effects of perceptual training on unmatched samples of brain-injured familial retarded children. *Journal of Abnormal Psychology,* 1965, *70,* 107–110.

FERNALD, G. M. Remedial techniques in basic school subjects. New York; McGraw-Hill, 1943.

FLAVELL, J. *The developmental psychology of Jean Piaget.* Princeton, New Jersey: D. Van Nostrand Co., 1963.

FROSTIG, M. *Frostig developmental test of visual perception.* Palo Alto: Consulting Psychologists Press, 1963.

FROSTIG, M., and HORNE, D. *The Frostig program for development of visual perception.* Chicago: Follett, 1964.

GATES, A. I. *Gates reading readiness test.* New York: Teacher's College Press, 1939.

GELB, A., and GOLDSTEIN, K. Psychologische Analysen hirnpathologischer Fälle. Berlin: Springer, 1920. Cited by Luria, A. R. *Human brain and psychological processes.* New York: Harper & Row, 1966b.

GELLERMANN, L. W. Form discrimination in chimpanzees and 2-year-old children. I. Discrimination of form per se; II. Form versus background. *Journal of Genetic Psychology,* 1933, *42,* 1–50.

GERSTMANN, J. Fingeragnosie. Wien. Klin. Wschr. 1924, *37.*

GETMAN, G. N. The visuomotor complex in the acquisition of learning skills. In J. Hellmuth (ed.), *Learning Disorders,* Vol. I. Seattle: Special Child Publications, 1965.

GETMAN, G., and HENDRICKSON, H. The needs of teachers for specialized information on the development of visuomotor skills in relation to academic performance. In W. Cruickshank (ed.) *The teacher of brain-injured children.* Syracuse: Syracuse University Press, 1966.

GIBSON, J. J. *The senses considered as perceptual systems.* Boston: Houghton Mifflin, 1966.

GIBSON, E. J., GIBSON, J. J., PICK, A. D., and OSSER, H. A developmental study of the discrimination of letter-like forms. *Journal of Comparative and Physiological Psychology,* 1962, *55,* 897–906.

GOLDSTEIN, K. *Language and language disorders.* New York: Grune & Stratton, 1948.

GOODENOUGH, F. L. *Draw-a-man test.* New York: Harcourt, Brace & World, 1926.

GRAHAM, F., and KENDALL, B. *Memory-for-designs test.* Missoula, Montana: Psychological Test Specialists, 1960.

GREGORY, R. L. *Eye and brain: The psychology of seeing.* New York: McGraw-Hill, 1966.

HALGREN, V. Rotation errors made by brain-injured and familial children on two visual-motor tests. *American Journal of Mental Deficiency.* 1955, *59,* 485–489.

HEAD, H. *Aphasia and kindred disorders of speech.* 3 vols. London: Cambridge University Press, 1926.

HÉCAEN, H., and AJURIAGUERRA, J. *Méconnaissances et hallucinations corporelles.* Paris: Masson, 1952.

HÉCAEN, H., and AJURIAGUERRA, J. Agnosie visuelle pour les objects inanimés par la lesion unilaterale gauche. *Rev. Neurology,* 1956, *94.*

HILDRETH, G. H., and GRIFFITHS, N. L. *Metropolitan readiness test.* Tarrytown, N.Y.: Harcourt, Brace & World, 1949.

HOLMES, G. Disturbances of vision by cerebral lesions. *British Journal of Ophthalmology,* 1919, *1.*

HOLMES, G. Cerebral integration of ocular movement. *British Medical Journal,* 1938, *2.*

HUNT, B., and PATTERSON, R. Performance of brain-injured and familial mentally deficient children on visual and auditory sequences. *American Journal of Mental Deficiency,* 1958, *63,* 72–80.

JACKSON, J. H. Case of large cerebral tumor without optic neuritis and with left hemiplegia and imperception. *Royal London Ophthalmic Hospital Reports,* 1876, Vol. VIII.

KAGAN, J., and LEMKIN, J. Form, color, and size in children's conceptual behavior. *Child Development,* 1961, *32,* 25–28.

KEPHART, N. C. *The slow learner in the classroom.* Columbus, Ohio: Charles E. Merrill, 1960.

KIRK, S. A. Hemispheric cerebral dominance and hemispheric equipotentiality. *Comparative Psychology Monographs,* 1935, *11* (5), Ser. No. 55.

KIRK, S. A. Extra-striate functions in the discrimination of complex visual patterns. *Journal of Comparative Psychology,* 1936, *21,* 145–160.

KIRK, S., and MCCARTHY, J. *Illinois test of psycholinguistic abilities,* Experimental ed. Urbana, Ill.: University of Illinois Press, 1961.

KIRK, S. A., MCCARTHY, J. J., and KIRK, W. D. *Illinois test of psycholinguistic abilities: Revised edition.* Urbana, Ill.: University of Illinois Press, 1968.

KLÜVER, H. Visual disturbances after cerebral lesion. *Psychological Bulletin,* 1927, *24.*

KLÜVER, H. An analysis of the effects of the removal of the occipital lobes in monkeys. *Journal of Psychology,* 1937, *2.*

KLÜVER, H. Visual functions after removal of the occipital lobes. *Journal of Psychology,* 1941, *11.*

KOLODNAYA, A. Y. The syndromes of spatial disturbances in brain wounds. *Uch. Zap. Mosk. University,* 1949, No. 111.

KOLODNAYA, A. Y. Disturbance of differentiation of right and left and the role of the cutaneous analyzer in its restoration. *Izvestiia,* Akademiia Pedagogicheskikh Nauk RSFSR, 1954, *53.*

KOPPITZ, E. *The Bender-Gestalt test for young children.* New York: Grune & Stratton, 1964.

LAMBETH, J. *What optometry and its related fields have to offer the reading teachers.* Duncan, Okla.: Optometric Extension Society, 1966.

LASHLEY, K. S. *The mechanisms of vision.* Vol. 1–XVII, Provincetown, Mass.: Journal Press, 1930–1942.

LING, B. C. Form discrimination as a learning cue in infants. *Comparative Psychological Monograph,* 1941, *17,* No. 2.

LISSAUER, H. Ein fall von seelenblindheit, nebst einem beitrag zur theorie derselben, *Arch. f. Psychiatry,* 1889–1890, *21,* 222–270.

LUBKER, B. J. The effect of training on cue-position patterning in discrimination problems. *Journal of Experimental Child Psychology,* 1964, *1,* 135–144.

LURIA, A. R. *Restoration of function after brain injury.* New York: Macmillion Co., 1963.

LURIA, A. R. *Higher cortical functions in man.* New York: Basic Books, 1966a.

LURIA, A. R. *Human brain and psychological processes.* New York: Harper & Row, 1966b.

LURIA, A. R., and HOMSKAYA, E. D. An objective study of ocular movements and their control. *Psychol. Beitr.* 1962, *6.*

LURIA, A. R., and SKORODUMOVA, A. V. The phenomenon of fixed hemianopsia. In *Collection of Memory of S. V. Kravkov,* Moscow, 1950.

McMURRAY, J. Visual perception in exogeneous and endogeneous mentally retarded children. *American Journal of Mental Deficiency,* 1954, *58,* 659–663.

METZGER, W. *Gestze des sehen.* Frankford on Main: Waldemar Kramer, 1953.

MISHKIN, M. Visual discrimination performance following partial ablation of the temporal lobe. *Journal of Comparative and Physiological Psychology,* 1954, *47.*

MISHKIN, M. Effects of small frontal lesion on delayed alternation in monkeys. *Journal of Neurophysiology,* 1957, *20.*

MISHKIN, M., and PRIBRAM, K. Visual discrimination performance following partial ablation of the temporal lobe. *Journal of Comparative and Physiological Psychology,* 1954, *48.*

MONEY, J. *Reading disability.* Baltimore: The Johns Hopkins Press, 1962.

MONTESSORI, M. *The Montessori method.* New York: Schocken Books, 1964.

MOORE, R., and GOLDIAMOND, I. Errorless establishment of visual discrimination using fading procedures. *Journal of Experimental Analysis of Behavior,* 1964, *7,* 269–272.

MORGAN, C. T., and KING, R. A. *Introduction to psychology.* New York: McGraw-Hill, 1966.

MUNK, H. *Uber die funktionen der grosshirnrinde.* Berlin: Hirschwaldt, 1881.

MUNN, N. L. *The evolution and growth of human behavior.* (2d ed.) Boston, Mass.: Houghton Mifflin Co., 1965.

MUNN, N. L., and STEINING, B. R. The relative efficacy of form and background in a child's discrimination of visual patterns. *Journal of Genetic Psychology,* 1931, *39,* 73–90.

MURPHY, HELEN A., and DURRELL, D. D. *Murphy-Durrell reading readiness analysis.* Tarrytown, New York: Harcourt, Brace & World, 1965.

NIELSEN, J. M. *Agnosia, apraxia, aphasia: their value in cerebral localization.* (2d ed.) New York: Hafner, 1962.

PAINTER, G. The effect of a rhythmic and sensory motor activity program on perceptual motor spatial abilities of kindergarten children. *Exceptional Children,* 1966, *33,* 113–116.

PAVLOV, I. P. *Complete collected works,* 6 Vols. Moscow: Izd. AN SSSR, 1949.

PENFIELD, W. *The excitable cortex in conscious man.* Springfield: Charles C Thomas, 1958.

PENFIELD, W., and JASPER, H. *Epilepsy and the functional anatomy of the human brain.* Boston: Little, Brown, and Co., 1954.

PENFIELD, W., and ROBERTS, L. *Speech and brain-mechanisms.* Princeton, N.J.: Princeton University Press, 1959.

PIAGET, J. *La naissance de l'intelligence de l'enfant.* Neuchâtel, Paris: Delachaux & Niestle, 1935.

POPPELREUTER, W. *Die psychischen Schadigungen durch Kopfschuss.* 2 vols., Leipzig: Voss, 1917–1918.

PRIBRAM, K. H. On the neurology of thinking. *Behavorial Science,* 1959, *4.*

PRIBRAM, K. H. A review of theory in physiological psychology. *Annual Review in Psychology,* 1960, *11.*

RAVEN, J. C. *Progressive matrices.* New York: Psychological Corp., 1956.

RHOS, L. A., RATLIFF, F., CONNSWEET, J. C., and CONNSWEET, T. The disappearance of steadily fixated visual test objects. *Journal of the Optometric Society of America,* 1953, *43,* 495–501.

ROACH, E. G., and KEPHART, N. C. *Purdue perceptual motor survey.* Columbus, Ohio: Charles E. Merrill Books, 1966.

ROSEN, C. An experimental study of visual perceptual training and reading achievement in first grade. *Perceptual and Motor Skills,* 1966, *22,* 979–986.

RUBIN, E. *Visuell wahrgenommene figuren.* Copenhagen: Glydendalske, 1921.

SEGUIN, E. *Idiocy and its treatment by the physiological method.* Albany: Columbia University, Brandow Printing Co., 1907 (first published in 1846).

SHEMYAKIN, F. N. The psychology of spatial ideas. *Scientific Reports of the State Institute of Psychology,* Moscow, 1940.

SHEMYAKIN, F. N. Investigation of topographical ideas. *Izvestia,* Akademiia Pedagogicheskikh RSFSR, 1954, 53.

SHEPARD, W. O. Learning set in preschool children. *Journal of Comparative and Physiological Psychology,* 1957, *50,* 15–17.

SHIF, Z. I. Disturbances of comparison in reading. *Izvestia,* Akademiia Pedagogicheskikh Nauk RSFSR, 1945, *2.*

STAATS, A., and STAATS, C. *Complex human behavior.* New York: Holt, Rinehart & Winston, 1963.

STRAUSS, A. A., and LEHTINEN, L. W. *Psychopathology and education of the brain-injured child.* New York: Grune & Stratton, 1947.

TAYLOR, E. A., and SOLON, H. A. *The fundamental readiness questionnaire and maladjustment.* New York: Reading and Study Skills Center, 1957.

TERMAN, L. M., and MERRILL, M. A. *Stanford-Binet Intelligence Scale,* Form L–M. Geneva, Ill.: Houghton Mifflin, 1960.

TEUBER, H. L. Perception. In J. Field, H. W. Magoun, and V. E. Hall (eds.) *Handbook of Physiology.* Section 1. Neurophysiology, Vol. 3. Washington, D.C.: American Physiological Society, 1960a.

TEUBER, H. L. *Visual field defects after penetrating missile wounds of the brain.* Cambridge: Harvard University Press, 1960b.

TEUBER, H. L. Effect of brain wounds implicating right or left hemisphere in man. In V. B. Mountcastle (ed.), *Interhemispheric relations and cerebral dominance.* Baltimore: Johns Hopkins Press, 1962.

TEUBER, H. L., and WEINSTEIN, S. Ability to discover hidden figures after cerebral lesions. *AMA Archives of Neurology and Psychiatry,* 1956, *76.*

WALDENFELSHEIM, J. S. Educational rehabilitation and visual education: an integrated approach. *The Optical Journal and Review of Optometry,* May 1962.

Webster's seventh new collegiate dictionary. Springfield, Mass.: G. & C. Merriam Co., 1963.

WECHSLER, D. *Wechsler intelligence scale for children.* New York: Psychological Corp., 1949.
WECHSLER, D. *Wechsler preschool and primary scale of intelligence.* New York: Psychological Corp., 1967.
WEDELL, K. Variations in perceptual ability among types of cerebral palsy. *Cerebral Palsy Bulletin,* 1960, 146–157.
WELCH, L. The development of discrimination of form and area. *Journal of Psychology,* 1939, *7,* 37–54.
YARBUS, A. L. The perception of an immobile graticule. *Bio fizika* I., 1956.

CHAPTER 4

HAPTIC PROCESSING

The human body provides several sensory avenues through which information can be obtained about the environment as well as the state of the body itself. Information may be obtained through the senses of hearing, sight, taste, smell, touch, and body movement. While the average person uses all sensory avenues, the auditory and visual systems seem to be the major input systems for acquiring information. In the absence of sensory impairment or excessive stimulation, sight and sound seem to dominate sensory awareness.

Children who can hear and see sometimes demonstrate a preference for using one system more than other systems. Children who are blind depend chiefly upon audition, while children who are deaf rely heavily upon vision. Touch and body movement are incorporated in educational programs for both the deaf and for the blind. Unfortunately, touch and body movement represent the main avenues of informational input for those children who are both deaf and blind.

While a great deal of information pertaining to the auditory and visual input channels is available, there is comparatively little information concerning the processing of cutaneous (touch) and kinesthetic (body movement) information. Also, the information which is available is widely scattered in various medical, child development, and psychological journals. Comparatively little information is found in most educational journals.

This situation is particularly distressing because experience in teaching deaf, blind, and deaf-blind children has clearly demonstrated the usefulness in using touch and body movement as a means of conveying information to children. Only a few teaching methods have emphasized touch (tactile) and body movement for teaching seeing and hearing children. Fernald's (1939) method of teaching remedial reading, writing, and spelling is a classic pattern of tactile-kinesthetic tracing technique. Gillingham and Stillman (1960), Straus and Lehtinen (1947) and Montessori (1912) have also developed tactile-kinesthetic methods of teaching. There is need, however, to investigate the potential value of touch and movement for teaching children who can hear and see. Similarly, there is need to learn more with respect to dysfunctions in the processing of cutaneous and kinesthetic information.

The purposes of this chapter are to discuss: (*a*) the central processing mechanism for obtaining information from the cutaneous and kinesthetic systems as they operate independently and simultaneously; (*b*) the kinds of sensory information which can be acquired; (*c*) the nature of dysfunctions in this processing system; (*d*) the procedures for assessment and treatment; and (*e*) the future research needs.

The Haptic Processing Mechanism

It is important to distinguish between vital sensations, gnostic sensations and haptic sensations. Vital sensations are largely unconscious and have to do with the automatic regulation of visceral events. Gnostic sensations are those such as vibration and position which require analysis and interpretation. For the purpose of this report, "haptic processing" refers to the integration of cutaneous and kinesthetic information. "Haptic perception" willl be used to denote information which is acquired as a result of the central processing and synthesis of cutaneous and kinesthetic information.

According to Gibson (1966) the haptic system is composed of different subsystems: (*a*) cutaneous touch gives perceptions of the skin and deeper tissue without movement; (*b*) touch-temperature refers to the combination of skin stimulation and vascular dilation or constriction; (*c*) touch-pain refers to the registration of pain; (*d*) haptic touch is the movement of the joints along with stimulation of the skin and deeper tissues; and (*e*) dynamic touch is muscular exertion in combination with stimulation of both the skin and the joints. Gibson also refers to "oriented touch" which is the continual feedback from the inner ear about the body's relationship with gravity and the ground.

881-152 O—70——4



Figure 4.—The Haptic System

The haptic system is activated when external stimuli come in contact with the skin and when muscular or skeletal movement occurs (see fig. 4). It should be noted that stimulation might be imposed on a passive subject or stimulation might be self initiated. Regardless of the active or passive nature of the stimulation, sense organs in the body are activated. These sense organs are named "mechanoreceptors."

According to Gibson (1966) mechanoreceptor cells are located throughout the body in the: (*a*) skin and deeper underlying tissue; (*b*) muscles and tendons which attach muscles to bone; (*c*) skeletal joints and connecting ligaments between all moveable bones; (*d*) blood vessels; and (*e*) hair cells located in the semicircular canals, utricle, saccule, and cochlea of the inner ear. These receptors convert the mechanical energy created by the stimulus into electrical energy which is transmitted along the neural pathways to the brain (see fig. 4).

The sensorimotor region of the brain, which is located in the parietal area or postcentral gyrus, serves as the analyzer for cutaneous-kinesthetic sensory information. The simultaneous input of cutaneous and kinesthetic information is integrated into haptic information or haptic perceptions about the body, its environment, and their interrelationships.

There is need to learn more about the nature and functioning of the haptic processing system and its various subsystems. Gibson (1966) highlights this need: "More than any other perceptual system, the haptic

apparatus incorporates receptors that are distributed all over the body and this diversity of anatomy makes it hard to understand the unity of function that nevertheless exists." (p. 134)

Kinds of Information Obtained by the Haptic System

Most school tasks, as do most acts in everyday life, require both touch and movement. The haptic system is important for obtaining information about object qualities, bodily movement, and their interrelationships. The simultaneous input of cutaneous and kinesthetic preception not only provides more informational input than either system alone, but by combining movement and touch it is possible to physically obtain various combinations of stimulus cues which would not be encountered if each system were used successively or independently of the other.

The haptic system provides two major kinds of information. The first category includes information about the environment such as: (*a*) geometric information concerning surface area or size, shapes, lines, and angles; (*b*) surface texture; (*c*) qualities of consistency such as hard, soft, resilient, or viscous; (*d*) pain; (*e*) temperature; and (*f*) pressure.

In the second category, bodily movement provides information about the body itself such as: (*a*) dynamic movement patterns of the trunk, arms, legs, mandible, and tongue; (*b*) static limb positions or postures; and (*c*) sensitivity to the direction of linear and rotary movement of the skull, limbs, and entire body. Body movement also provides information about the location of objects in relation to the body itself.

Duration of body movement through space supplies information about the areas where objects are not encountered. This is accomplished by physically seeking out and contacting the concrete boundaries of an area.

Muscular effort and tension or strain in the joints, ligaments, and tendons provide information about weight or resistance to gravity. Pushing and pulling an object in the horizontal direction or lifting and lowering objects in the vertical direction give information about the relative weights of objects. Muscular effort and strain also supply information about the relationship of one's own body to gravitational pull.

Description, Assessment, and Treatment of Cutaneous Processing Dysfunctions

As early as 1874, Wernicke reported case studies in which pain and tactile sensation remained intact, but in which complex discriminatory sensation was affected. Since that time several cutaneous dysfunctions have been identified. These include failure to: (*a*) identify the presence of pressure on the skin; (*b*) localize the point of mechanical stimulation; (*c*) differentiate two or more stimuli which are applied simultaneously; (*d*) indicate the direction of an object moving over the surface of the skin; and (*e*) register sensitivity to pain and temperature.

A number of case studies have been reported which have shown that the synthesis of tactile stimuli may be disturbed by lesions in the parietal areas of the cerebral cortex (Luria, 1966). Lesions in the posterior central gyrus region on one side of the body were found to be accompanied by absence of cutaneous sensation on the opposite side of the body (Head, 1920). Semmes, Weinstein, Ghent, and Teuber (1960) also found that lesions in the sensori-motor or postcentral region of the left hemisphere may reduce tactile sensation in the right upper limb. Their studies suggest that the cerebral localization of the higher tactile functioning is more highly concentrated in the dominant hemisphere. The significance of these areas is further emphasized by Luria (1966) who reports that lesions outside the sensori-motor postcentral and posterior parietal regions of the left hemisphere usually cause no disturbance of tactile sensation.

Single-Point Discrimination and Localization (Touch versus No Touch)

A common procedure in the investigation of tactile functioning is to exclude sensory input from the visual and/or kinesthetic receptors by screening and immobilizing the limb. The threshold of tactile stimulation can be studied by applying tactile stimuli to the subject's fingers, palm, forearm, and shoulder with increasing pressure. The frequency intervals with which stimuli are applied are varied to allow for the aftereffects of sensation and fatigue. When a point stimulation increases in intensity, the cutaneous deformation becomes enlarged and deeper and a greater number of receptors are activated.

Gibson (1966) mentions several instruments which are used to create mechanical energy, mechanically displacing or deforming the skin, and activating the mechanoreceptors in the skin. These stimuli include a rounded wedge for rubbing, a sharp wedge for scraping, a small cylinder for rolling, and tufts of hair for brushing. Each of these instruments can be used to apply different kinds of pressure. Torsion of the skin can be produced by twisting a rubber stimulator clockwise or counterclockwise to see if the subject can indi-

cate the direction of the torsion. Forceps are used to test sensitivity to stretching or pinching. Sensitivity to motion over the surface of the skin, lateral traction, is measured by drawing a long rod over the skin region (Gibson, 1962). Thermal sensitivity can be tested by applying metallic objects which have been heated or cooled to different temperatures.

Several studies have used surface texture to assess tactile discrimination in the absence of vision. Rough and smooth paper, commercial grades of sandpaper (Stevens and Harris, 1962), and fleece and untwisted rope fibers (Binns, 1937) were used to study tactile perceptiveness. It is interesting that little tactile information was obtained by mere touching. In order to discriminate, it was necessary to rub the stimulus object or pull the stimulus object through the fingers. Mechanical friction seemed necessary in order to obtain sufficient information about subtle differences in surface texture so that comparative judgments could be made.

Classifying the intensity of two, three, or more stimuli which have been applied in random order requires greater discrimination than does differentiating between touch and no touch. Diminution of sensitivity on one side of the body usually indicates lesions in postcentral divisions of the contralateral hemisphere or of the corresponding conductive tracts.

Tests for tactile localization can be administered by simply having the subject indicate the part of the skin touched by the instrument. The task can be made more difficult by asking the subject to point to the corresponding point on the opposite limb. This requires the subject to preserve the tactile localization and then to identify the symmetrically opposite area of the skin. Error can be measured in millimeters or centimeters.

Sensitivity to pain is the consequence of excessive or intense mechanical, thermal, electrical, or chemical stimulation. According to Geldard (1953) pain is usually sharply localized and tends to elicit prompt withdrawal reactions. Pain has value as a carrier of useful information because it produces withdrawal reactions to environmental conditions which may be injurious. Contact with an object in the environment, such as a pin or a hot stove, is often the only way a child is able to learn whether a physical contact is injurious or not, Sternbach (1963, 1968) reviews the clinical and experimental literature on pain and concludes that otherwise "normal" individuals are apparently insensitive to pain, yet have acquired all the expected social behaviors without the assumed negative reinforcement, assumed necessary to learning such avoidance behaviors, such as avoiding fires and sharp objects. Great individual differences in sensitivity to and awareness of pain are found in randomly-sampled populations. Gibson (1966) points out the need to learn more about the nature of pain and how it is perceived.

Multiple-Point Discrimination and Tactile Recognition

The touch compass test is useful in investigating the threshold at which the subject is able to differentiate between one-point and two-point stimuli. Two points are placed in contact with the skin as closely as possible and then gradually separated until the subject is aware of two-point sensation. In cases where the parietal region has suffered damage, the subject often has difficulty perceiving when two stimuli are applied simultaneously to the skin (Head, 1920; Bender and Teuber, 1947, 1948; and Bender, 1952).

The two points of the compass can be presented simultaneously or successively. For example, Teuber (1959) found that subjects with lesions of the parietal cortical region are able to distinguish tactile stimuli on both the right and left sides of the body, provided the stimuli are applied to the skin surface separately. In the event one stimulus is applied to the right side of the body and a second stimulus is applied, simultaneously, to the left side of the body, the individual with damage to the right hemisphere will notice only the stimulus on the right side. He will fail to notice the stimulus on the left side. This phenomenon represents a breakdown in the discrimination of multiple-point stimuli.

The movement of an object across the skin's surface triggers a temporal sequence of successive sensations. Failure to process this tactile information results in failure to indicate the direction of the object moving across the skin's surface. This phenomenon represents a dysfunction in synthesizing successive multiple-point stimulation. An extensive discussion of the skin senses has been conducted by Kenshalo (1968).

Procedures for assessing the processing of complex cutaneous information have been developed by Strauss and Werner (1938), Strauss and Lehtinen (1947), Hauessermann (1958), Benton (1959), Kinsbourne and Warrington (1963–1964), and others. Hauessermann's (1958) tests of tactual sensitivity provide a basis for the diagnosis of tactile agnosia in children from the ages of 4 to 6 years. The tests determine if the subject is able to use tactile scanning for the perception of slight differences in texture, and the recognition of familiar objects by touching their form and texture without seeing them.

Kinsbourne and Warrington (1963–1964) used three separate tests to assess the ability to perform discriminations based on the classification of sensory input from the fingers in terms of the individual fingers which were stimulated in their correct spatial arrangement. Two of the tests require verbal responses. The subjects ranged in age from 4½ years to 7½ years. Results of their tests indicated that the acquisition of "finger sense" is a maturational process. The ability to differentiate sensory input from the fingers was acquired by the approximate age of 5½ years in 50 percent of their group of normal children. Kinsbourne and Warrington's procedures for evaluating "finger sense" are described as follows—

A. Finger Differentiation Test: In the finger differentiation test, the subject faced the examiner with his eyes closed. One hand was positioned with the palm down and the fingers spread. Two points were touched simultaneously on the fingers, sometimes both on the same finger and sometimes on two adjacent fingers. The examiner then asked the subject how many fingers he was touching. He maintained contact until an answer was given by the subject and long enough after the answer to permit the subject to verify the answer under direct vision.

B. The "In Between" Test: In the "in between" test the subject's hand was positioned with the palm down and the fingers spread. The examiner touched two fingers simultaneously and asked the subject to tell how many fingers were in between the ones he was touching. The examiner maintained contact until answered and long enough to permit the subject to verify the answer under direct vision.

C. The Finger Block Test: Four wooden blocks with distinctive shapes were utilized in the finger block test. The subject's fingers were molded around one of four corresponding test blocks while his eyes were closed. The subject was then directed to open his eyes and without looking at the block in his hand, pick out the corresponding one on the table. After making his choice, the subject was allowed to verify his answer by direct visual comparison.

Recognizing objects by touch and distinguishing shapes, letters or numbers drawn on the skin require integration of sensations from a series of points on the skin's surface. Wernicke (1874) and Nielsen (1946) believed that failure to recognize objects and shapes represents a symbolic disorder. In contrast, Bay (1944) and others have viewed tactile processing dysfunctions as more elementary sensory disorders which may manifest themselves as the tactile task becomes more complex. Failure in object recognition may be due to a basic sensory dysfunction in synthesizing complex cutaneous information (Luria, 1966). There is need to question, study and clarify the traditional distinctions which have been drawn with respect to higher symbolic tasks and complex sensory integrative tasks.

The lack of information about cutaneous processing suggests the need for further research in this area. Anderson (1966) suggested several possible research directions: (1) isolation of research variables relating touch to learning; (2) increased work with clinical populations; (3) study of developmental trends in touch perception with clinically defined groups; (4) exploration of the extent to which enrichment or deprivation of tactile scanning experiences influences the development of symbolic skills; and (5) selection of diagnostic data which is more directly related to training procedures.

Description, Assessment, and Treatment of Kinesthetic Processing Dysfunctions

The kinesthetic sense makes little demand on our attention. Yet, simple acts such as lifting the foot from the ground, reaching for an object, or sitting down are continuously monitored by incoming signals from the mechanoreceptors. According to Wyburn, Pickford, and Hirst (1964), even without seeing movement we are aware of positions taken by different parts of the body. Tissue deformations, tensions, flexion and extension stretching, compressions, and change in length of muscle or ligaments all generate kinesthetic information which aid in controlling movement. The positioning and movement of the joints, muscular contraction and tension resistance of the tendons trigger kinesthetic feedback. Breakdowns in kinesthetic processing can interfere with the feedback of several different kinds of kinesthetic information.

Joint and Muscle Sensitivity to Movement

The importance of joint sensitivity to the perception of space and movement has not received sufficient attention. Contrary to widespread belief, "* * * muscle sensitivity is irrelevant for the perception of space and movement, whereas joint sensitivity is very important for it. In short, we detect the angles of our joints, not the length of our muscles" (Gibson, 196, p. 109).

The amount of information which is obtained from muscles and tendons is limited. Nerve endings in the muscles and tendons are triggered by tension. The muscle fibers register the stretch of fiber, and the tendon receptors register strain. Resistance to gravity is obtained through muscular effort. Since muscle

length is not correlated with muscle effort, the muscle itself provides comparatively little information. Muscular effort in combination with mechanoreceptors in the skin, joints, ligaments, and tendons yields the most information.

In addition to the function of permitting mobility of the articulated bones, the relative position and movement of the bones is registered by the receptors in the ligaments and the receptors in the capsules of the joints. Consequently, awareness of position and movement of the joints depends solely on the receptors in the joints themselves. Reference to some mysterious "muscle sense" to explain kinesthetic sensation is unnecessary and runs contrary to all known facts concerning the muscle-stretch receptors (Rose and Mountcastle, 1959).

The joint receptors discharge at a given rate for a given angle of a joint. When the angle changes, the rate of discharge changes. Sensitivity to these changes was demonstrated in studies by Goldscheider (1898) and Geldard (1953), who found that subjects could detect the bending of a single joint as little as a fraction of 1°. The angle of one joint has little meaning, however, unless it is related to the angles of all mobile joints in the body of which there are approximately one hundred. The angular position of every bone of the body is articulated with the body frame which is anchored in the direction of gravity pull. The sensory feedback from angle changes and rotations of the joints provide perception of space and movement, particularly in the absence of vision. Motor behavior consists, in part, of a succession of angular changes which give the impression of continuous sensory feedback. Troland (1929) has described movement as a succession of many different postures.

All five fingers of each hand are frequently used to provide sensory information (Kohler and Dinnerstein, 1949). Finger span, the dimension in space between the thumb and index finger, is often a source of information concerning the dimensions of small objects (Katz and MacLeod, 1949; Kelvin, 1954). The angular position of the bones in the fingers is also used to help determine geometrical information such as the shape of surfaces and the arrangement of objects in space. Similarly, sensory information about the size of large objects may be obtained by extending both arms, thus rotating shoulder, elbow, and wrist joints, thereby determining kinesthetically the span or distance between the hands. There is need to develop procedures for training increased joint sensitivity to joint and muscular movements.

Linear and Rotary Movement

Kinesthetic information which is registered by the inner ear includes sensitivity to linear movement, acceleration, deceleration, directional changes, falling, and clockwise or counterclockwise rotary motion or spin. Information is obtained about the vertical and horizontal axes of the world, the gravitational point of reference and an awareness of the posture of the body. The vestibular apparatus in the inner ear registers stops, starts and changes in direction but it is not sensitive to uniform movement in a constant (straight-line) direction. The vestibular apparatus is defined as the utricle with the three semicircular canals and the saccule. These enclosures are filled with fluid into which the hairs of the receptors penetrate. The fluid moves with movement of the body and the receptors are stimulated by the motion of the fluid.

Movement Patterns and Static Postures

For voluntary controlled movement to occur, it is necessary to receive continuous feedback from the kinesthetic system. The mechanoreceptors which are distributed around the ligaments, joints, and deep tissue normally provide this feedback (Gibson, 1966).

Lesions in the postcentral divisions of the cerebral cortex have been found to be related to motor disturbances in which the power of the muscle remained unimpaired but voluntary movements could not be performed (Foerster, 1936). The motor impulses lost their selectivity and produced simultaneous contraction of both the agonistic and the antagonistic muscles. In less severe cases, subjects experience difficulty in reproducing finger position. Foerster also found similar problems arising from damage to the parietal area.

Recent thinking has emphasized the importance of the preparatory process for motor behavior. Before a voluntary motor act can be executed, cortical areas in the sensori-motor region (postcentral and precentral regions) must simultaneously organize the movements in the system of external spatial coordinates and analyze impulses arriving from the muscles and joints. During the early stages of motor development, both kinesthetic and optic afferent impulses play an important role in developing voluntary movement (Zaporozhets, 1960). As further development takes place, however, language is used as a mediator for planning voluntary action and kinesthetic and optic afferent components of movement assume a supporting role (Bernshtein, 1947).

Head (1920) and Denny-Brown (1958) found that a lesion of the posterior divisions of the sensori-motor region (area 4) results in disturbances of spatial inte-

gration or to kinesthetic schemes of movement upon which motor behavior is based. Motor impulses lose selectivity and go to agonists and antagonists at the same time.

The work of Piaget (1935) and others shows that kinesthetic visual and vestibular analyzers are responsible for processes of inspection, palpation, head orientation, eye movement, and the development of spatial relationships. "Before a voluntary motor movement can be carried out, the visual, vestibular, or acoustic impulses must first be recoded into a definite system of kinesthetic signals." (Luria, 1966, p. 174) The kinesthetic basis of movement has been placed in the postcentral divisions of the cortical nucleus of the motor analyzer. Exteroceptive signals are recoded into kinesthetic combinations and impulses integrated into simultaneous groups.

Failure to process information about movement patterns can be investigated by having the subject reproduce various movement patterns or postures with his limbs. Before administering such a test, however, a neurological examination should be given to provide information about manifestations of muscle power, accuracy of movement, disturbances of muscle tone, ataxia, and hyperkinesis.

Luria (1966) describes a procedure for assessing optic-kinesthetic organization of a complex movement. The subject is asked to reproduce different finger positions demonstrated by the experimenter. Visual input is blocked by placing the subject's hand through a hole in a screen. Failure on this test is noted by diffuse movements or the inability to achieve the necessary selection of movement. Mirror-image reversals of movements are not necessarily connected with kinesthetic disturbances. They may be eliminated if the examiner either sits beside the subject or asks the subject to reproduce with his left hand, movements presented by the right hand. The tendency to perseverate a movement may be an indication of a lesion of the anterior division of the cerebral cortex. Pathological inertia of motor acts accompanied by postural praxis is usually a sign of a lesion (Luria, 1966).

The examiner may vary the test by positioning one of the patient's hands and asking him to reproduce the position with the other hand while keeping his eyes shut, or asking the subject to execute a movement from a verbal command. This removes the optic afferent component of the motor analyzer. Luria (1966) also discusses tests which help identify defects in the optic-spatial organization of the motor act.

Several tests help identify the presence of defects in the kinesthetic basis for accomplishing hand movements. The examiner places the subject's hand or fingers in a certain angle or position and asks him to reproduce this angle with the same hand and with the opposite hand. Then the examiner asks the subject to shut his eyes and reproduce the angle with the same hand and with the opposite hand. Poor performance on these tasks or the gradual increase of errors as fatigue develops suggests that there is a deficit in kinesthetic analysis in the opposite hemisphere.

There are a number of procedures which are used to assess the functioning of deep muscle and joint sensation. The subject usually sits upright, with his eyes covered, and his arm, hand, and fingers in a fixed position. The examiner then moves the subject's arm, hand, or fingers up, down, and to the side. The subject is asked to indicate the direction of movement. Subjects are also asked to reproduce the position of a limb independently, or move the opposite limb to a corresponding position. The angle of the forearm may be moved passively from one angle to another to determine if the subject is able to: (*a*) Determine if the two movements were the same or different; (*b*) repeat those consecutive movements; or (*c*) reproduce the movements with the other limb.

Disturbances in deep muscle and joint sensations, particularly in the upper limb, tongue and lips have been found to arise from lesions in the postcentral and posterior parietal regions of the cortex of the opposite hemisphere. Disturbance of kinesthetic sensation is found mainly in the contralateral upper limb. These conditions lead to evaluative errors.

The reproduction of actions may be severely impaired when the object is not visibly present. When the object is present, vision may help support the subject's attempt to reproduce actions. This suggests that the kinesthetic organization of action is a basic element of symbolic actions.

The disturbance of the kinesthetic basis of speech has been classified as "afferent kinesthetic motor aphasia." Disruption of the coordinated movements essential for the kinesthetic basis of motor acts results in failure to organize and execute articulation in speech (Liepmann, 1913).

As early as 1861, Broca (1861a, 1861b) described motor aphasia for the motor images of words as distinct from all other sensory and motor disorders. Further, he localized this disorder within the third frontal gyrus. Nissl von Meyendorff (1930) found that motor aphasia also resulted from lesions in the Rolandic area and in-

volved the inferior portions of the postcentral kinesthetic region of the cortex as well.

Lesions of the inferior divisions of the postcentral region of the left hemisphere have also been found to result in apraxia of the tongue, lips, and palate (Luria, 1966). While the subject can produce sounds, he is unable to select the correct positions of lips, tongue, or palate. This requires concentration and results in frequent sound substitutions while speaking because the subject is unable to assume the correct position with the articulators in transferring from one sound to the next. In contrast, the condition of dysarthria is characterized by the inability to produce individual sounds and by slurred speech and monotonous speech patterns. In some cases entire phrases may be produced easily while great difficulty is experienced in articulating specific sounds. Such abnormalities of speech should be followed up by an investigation of the articulatory mechanism and how it functions.

Many tests for oral praxis usually require the subject to reproduce movements of lips, tongue, and face, such as stretching lips; baring teeth; extending tongue flatly; folding it up; puffing out cheeks; and, placing tongue between teeth. While the subject is performing these acts, his lips, tongue, palate, and facial muscles (controlled by the inferior portion of sensori-motor zone and adjacent parieto-temporal areas) should be observed for symptoms such as protruding lips; asymmetry of movement when showing teeth, puffing cheeks, wrinkling the brow, frowning, squinting, or positioning the tongue; limited range of tongue movement; excesive salivation; smooth movement substituted by tremors, spasms, or tension; and paresis of the soft palate or facial muscles. These all represent peripheral disorders of the articulatory act.

The subject is asked to reproduce two or three movements in succession either from demonstration or verbal command. These may be the same or different movements. Rapid performance of the first movement followed by the inability to switch movements or a fixation on one movement may be indicative of a lesion in the anterior divisions of the motor cortex. The examiner may gain further information by contrasting the subject's natural performance of an action with his performance upon request. The inability to act on command, while able to function in the real situation is characteristic of those with brain lesions.

Writing and reading activities are often affected by the kinesthetic form of motor aphasia. A study by Nazarova (1952) demonstrated that when first and second grade pupils were not permitted to articulate, the number of errors in writing increased five to six times. Children use articulatory movement as a mediator to analyze the sound composition of words during the initial stages of learning how to read and write. Thus, difficulty in organizing and producing speech sounds, whether or not due to lesions in the postcentral cortical divisions, may interfere with learning how to write. The inability to organize and produce appropriate sounds may interfere with reading, particularly in words which have complex sound sequences requiring rapid transitions. Sound substitutions and sound distortions resulting from failure to produce the correct auditory language signals for their graphic representations will interfere with comprehension of what s to be read.

Future research is needed to provide a more thorough description of these disorders and to develop more effective procedures for their assessment. Causes for failure to produce complex voluntary movement must be differentiated, such as: (*a*) faulty kinesthetic processing; (*b*) inadequate muscle power and tone; (*c*) dysfunctions in the optic-spatial afferent system; and (*d*) failure to inhibit one set of muscle groups while exciting other muscle groups.

There is need, also, to determine if training can improve the selection and organization of kinesthetic impulses for bringing about increased control over voluntary movement.

The literature reveals that special teaching procedures have been developed for training cutaneous-kinesthetic abilities in normal young children (Montessori, 1912; Scagliotta, 1967). Remedial taining in the tactual-kinesthetic mode has also been recognized by educators as a means of reaching children with special learning problems. Monroe (1932), Fernald (1939), Strauss and Lehtinen (1947), Gillingham and Stillman (1960), Kephart (1962), and Frstig (1964–1965) all suggest methods of remedial reading, writing, spelling, and arithmetic which employ cutaneous-kinesthetic training techniques. The Hegge, Iirk, and Kirk (1948) remedial reading drills uses a grahovocal method for teaching reading.

These procedures have been used for ameliortion of the basic disability as well as for improving the fnctioning of one modality to compensate for the dysfnction of a second modality. A major issue in trainig is whether or not haptic processing disabilities can b influenced by various forms of training exercise or whether it is necessary to work with the residual bilities and teach the subject to compensate by utiling his assets.

A review of the research reveals that there is ttle known about either the restoration or the compension

of haptic processing dysfunctions in children. It is very likely that the training programs devised for deaf, blind, and deaf-blind children have the greatest applicability for teaching hearing and seeing children who have difficulty processing and integrating cutaneous-kinesthetic information.

Further research should investigate the amelioration of haptic processing dysfunctions, as well as the ways in which the haptic modality can be used to reinforce the learning of children who have other kinds of disorders. Increased attention should be directed toward haptic input as a compensatory source of information for children whose auditory or visual processing mechanisms are impaired.

Directions for Future Research

There is need for research on the processing and integration of cutaneous and kinesthetic information. In addition to acquiring more precise behavioral descriptions of the dysfunctions in the haptic system, there is need to: (*a*) develop more systematic assessment procedures for describing deviant behaviors; (*b*) devise more effective diagnostic procedures for linking specific behavioral symptoms with the causal factors; and (*c*) study the effectiveness of different treatment procedures with specific dysfunctions.

Most of the literature has been concerned with utilizing the cutaneous-kinesthetic processing to reinforce other sensory modalities which may be defective in children. A major issue which needs to be resolved is whether or not it is possible to improve tactile-kinesthetic abilities in children whose tactile-kinesthetic sensory modality is distorted or deficient. Research is needed, also, to explore the compensatory advantages of using the haptic modality to compensate for children who have dysfunctions in auditory or visual processing.

REFERENCES

Anderson, R. P. Physiologic considerations in learning: The tactual mode. In J. Hellmuth (ed.) *Learning Disorders.* Vol. 2. Seattle: Special Child Publications, 1966.

Bay, E. Zum Problem der taktilen Agnosie. *Deutsche Z. Nervenh.*, 1944, 156.

Bender, M. B. *Disorders of perception.* Springfield, Ill.: Charles C Thomas, 1952.

Bender, M. B., and Teuber, H. L. Spatial organization of visual perception following injury to the brain. *AMA Archives of Neurology and Psychiatry,* 1947, p. 58; 1948, p. 59.

Benton, A. L. *Right-left discrimination and finger localization.* New York: Hoeber and Harper, 1959.

Bernshtein, N. A. *The Construction of Movements.* Moscow: Medgiz, 1947.

Binns, H. Visual and tactual judgment as illustrated in a practical experiment. *British Journal of Psychology,* 1937, *27,* 404–410.

Broca, P. Perte de la parole etc. *Bull. Soc. Anthrop.*, 1861a, 2.

Broca, P. Remarques sur le siege de la faculte du langage articule. *Bull. Soc. Anthrop.*, 1861b, 6.

Denny-Brown, D. The nature of apraxia. *Journal of Nervous and Mental Diseases,* 1958, 126.

Fernald, G. *Remedial techniques in basic school subjects.* New York: McGraw-Hill, 1939.

Foerster, O. Symptomatologie der Erkrankungen des Gehirns. Motorische Felder und Bahnen. Sensible corticale Felder. In O. Bumke and O. Foerster (eds.) *Handbuch der Neurologie,* Vol. 6. Berlin: Springer, 1936.

Frostig, M. Corrective reading in the classroom. *The Reading Teacher,* 1964–1965, *18,* 573–580.

Geldard, F. A. *The human senses.* New York: Wiley, 1953.

Gibson, J. J. Observation on active touch. *Psychological Review,* 1962, *69,* 477–491.

Gibson, J. J. *The senses considered as perceptual systems.* Boston: Houghton Mifflin Co., 1966.

Gillingham, A., and Stillman, B. *Remedial training for children with specific disability in reading, spelling, and penmanship.* Cambridge, Mass.: Educators Publishing Service, 1960.

Goldscheider, A., *Gesammelte Abhandlungen von, A. Goldscheider* (2 vols.), Leipzig: Barth, 1898.

Hauessermann, E. *Developmental potential of preschool children.* New York: Grune & Stratton, 1958.

Head, H. *Studies in Neurology,* 2 vols. London: Oxford Medical Publishers, 1920.

Hegge, T. G., Kirk, S. A., and Kirk, W. D. *Remedial reading drills.* Ann Arbor, Mich.: George Wahr Publishing Co., 1948.

Katz, D., and MacLeod, R. B. The mandible principle in muscular action. *Acta Psychology,* 1949, *6,* 33–39.

Kelvin, R. P., Discrimination of size by sight and touch. *Quarterly Journal of Experimental Psychology,* 1954, 6, 23–34.

Kenshalo, D. R. *The skin senses.* Springfield, Ill.: Charles C Thomas, 1968.

Kephart, N. C. *The slow learner in the classroom.* Columbus, Ohio: Charles E. Merrill, 1962.

Kinsbourne, M., and Warrington, E. K. The development of finger differentiation. *Quarterly Journal of Experimental Psychology,* 1963–1964, *15–16,* 132–137.

Kohler, W., and Dinnerstein, D. Figural after-effects in kinesthesis. In *Miscellanea Psychologica Albert Michotte,* Louvain: Editions de l'Institut Superieur de Philosophie, 1949.

Liepmann, H. Motorische Aphasie und Apraxie. *Monatschr. Psychiat. u. Neurol.* 34, 1913.

Luria, A. R. *Human brain and psychological processes.* New York: Harper & Row, 1966.

Monroe, M. *Children who cannot read.* Chicago: University of Chicago Press, 1932.

Montessori, M. *The Montessori method.* New York: Frederick A. Stokes, 1912.

NAZAROVA, L. K. The role of speech kinesthesias in writing. *Sovet Pedag.* 1952, 6.

NIELSEN, J. M. *Agnosia, Apraxia, Aphasia.* Los Angeles, 1946.

PIAGET, J. La naissance de l'intelligence de l'enfant. Neuchatel, 1935.

ROSE, J. E., and MOUNTCASTLE, V. B. Touch and kinesthesis. Chapter 17 in *Handbook of Physiology.* Vol. 1, *Neurophysiology,* American Physiological Society, 1959.

SCAGLIOTTA, E. G. Tinker boards-A tactile kinesthetic approach to learning. *Exceptional Children,* 1967, *34,* 129–131.

SEMMES, J., WEINSTEIN, S., GHENT, L., and TEUBER, H. L. *Somatosensory changes after penetrating brain wounds in man.* Cambridge: Harvard University Press, 1960.

STERNBACH, R. A. Congenital insensitivity to pain: A critique. *Psychological Bulletin,* 1963, *60,* 252–264.

STERNBACH, R. A. *Pain: A psychophysiological analysis.* New York: Academic Press, 1968.

STEVENS, S. S., and HARRIS, J. R., The scaling of subjective roughness and smoothness. *Journal of Experimental Psychology,* 1962, *64,* 489–494.

STRAUSS, A., and LEHTINEN, L. E. *Psychopathology and education of the brain-injured child.* New York: Grune & Stratton, 1947.

STRAUSS, A. and WERNER, H. Deficiency in the finger schema in relation to arithmetic disability (finger agnosia and acalculia). *American Journal of Orthopsychiatry,* 1938, *8,* 719–725.

TEUBER, H. L. Some altercations in behaviour after cerebral lesion in man. In: Evolution of Nervous Control. American Association for the Advancement of Science. Washington, D.C., 1959.

TROLAND, L. T. *The principles of psychophysiology.* Vol. 1, *Problems of psychology and perception.* Van Nostrand, 1929.

VON MEVENDORFF, NISSL. *Vom Lokalisationsproblem der artikulierten Sprache.* Leipzig: Barth, 1930.

WERNICKE, C. *Der aphasische Symptomenkomplex.* Breslau: Cohn & Weigart, 1874.

WYBURN, G. M., PICKFORD, R. W., and HIRST, R. J. *Human senses and perception.* Canada: University of Toronto Press, 1964.

ZAPOROZHETS, A. V. Development of voluntary movement in the child. *Izd. Akad. Ped. Nauk* RSFSR; Moscow, 1960.

DYSFUNCTIONS IN THE SYNTHESIS OF SENSORY INFORMATION